

Dental Hygiene
Student Handbook
2023 – 2024



UTHealth[®] Houston

School of Dentistry

PREFACE

The University of Texas School of Dentistry at Houston (UTSD) Dental Hygiene Program Student Handbook should serve you well as a student in the dental hygiene program. It provides general and professional information as well as clinic information, clinical procedures, and program policies. You will want to keep the Handbook close at hand at all times. The Handbook is updated annually, but changes may occur in the interim and you will be informed of these. Use it as your guide and refer to it when appropriate.

In a desire to live by our philosophy, we aspire to be the model Dental Hygiene Program for the 21st century dedicated to advancing the health of the people of the State of Texas, the nation, and our global community through educating compassionate health care professionals and innovative scientists and through discovering and translating advances in the social and biomedical sciences to treat, cure, and prevent disease now and in the future". The UTSD Dental Hygiene Program's goals are as follows:

- Recruit well-qualified, diverse students and educate them to be qualified oral health care professionals in their chosen field of dental hygiene by preparing them for future practice in a highly-technologic world—amid a population that is aging, ethnically diverse and consumer-oriented.
- Graduate clinically competent students who can provide comprehensive patient care.
- Educate students to provide patient care in adherence to guidelines of a comprehensive quality assurance and risk management program.
- Provide service learning, research and community outreach experiences that enrich students' professional development and reinforce their clinical education.

As a UTSD student, you are encouraged to maintain close communication with your assigned facilitator, course directors, advisor, and other faculty in order to make your education as smooth as possible. If your faculty feels that you would benefit from counseling, they may recommend The University of Texas Health Science Center at Houston's (UTHealth) Student Health and Counseling Services or you may make an appointment on your own by calling 713-500-5171.

Counselors are available to provide counseling in regards to personal and/or academic issues. More information can be found on their website at <https://www.uth.edu/studenthealth/student-counseling-services/about.htm>.

Please refer to the UTSD Academic Catalog and the UTHealth General Information Catalog for more information about the Americans with Disability Act (ADA) if you feel that you may require any disability accommodations. The catalogs will also provide additional information on life at the UTHealth and the School of Dentistry.

On behalf of the dental hygiene faculty, I welcome you into the program and promise to support you in every possible way to ensure your academic success.

Victoria Patrounova, R.D.H., M.H.A.
Associate Professor and Program Director
Department of Periodontics & Dental Hygiene

TABLE OF CONTENTS

Section A - Student Information

Administration	7
Vision & Mission	7
General Guidelines	8
Dental Hygiene Community Liaison Council	8
Disability Accommodation	8
Procedure to File Complaints with The Commission on Dental Accreditation	9
Advisory System	10
Protocol for Resolution of Student Concerns	10
Dental Hygiene Complaint, Suggestions, & Compliments Form	10
Ethics and Professionalism	10
Dental Hygiene Licensure Eligibility	11
Academic Integrity	11
Professionalism	14
Corrective Action	14
Student Incident Report	15
Clinical Infractions	15
Cardinal Infractions	16
Attendance	16
Religious Holy Day	17
Severe Weather and School Closure	17
Procedures for Reporting Absences	17
ID Badges	17
Appearance Guidelines	18
Examinations	20
Academic Standards	20

Grade Grievance Procedure	21
Warning, Probation, and Dismissal	21
Academic Action and Appeal Process	22
Remediation	23
Student Financial Services Policy Regarding Satisfactory Progress	23
Student Withdrawal	23
Leave of Absence	23
Readmission	24
Academic Counseling	24
Clinical Equipment Information	24
Instrument Return/Replacement Policy	25
Solicitation of Patients	25
Use of Social Media	25
Clinical Patients for Licensure Examinations	25
<i>Section B – Professional Information</i>	
Oath of the American Dental Hygienists' Association	27
Student Chapter American Dental Hygienists' Association	27
The Dental Hygiene Program Pin	30
Code of Professional Ethics	30
Awards and Honors	31
UTHealth Endowments and Scholarships	34
<i>Section C – General Clinic Information</i>	
General Guidelines for Ergonomic Positioning Practice	37
Health History Guidelines/Medical Consultation	40
Prophylactic Antibiotic Therapy	41
Classifications of Hypertension	43
ASA Physical Status Classification System	45
Pregnancy	47
Blood Sugar Levels	47
Medical Consultation Record - Student Instructions	48
ABCDE System of Recognizing Melanoma	49
Gingival Description	49
Calculus Classification	50
Age Criteria	51
Special Need Patient	50
Classifications of Periodontal Health Diseases	51
Periodontal Chart	54
Caries Diagnostic Criteria	56
Occlusion-Patient Evaluation	57
Electronic Health Record	58

Supplemental Fluoride	62
General Appointment Sequence	62
Protocol for Topical Anesthetic	67
Protocol for Oraqix®	68
Protocol for Arestin®	69
Protocol for Local Anesthetic	70
Clinic Assistant Rotation	72
Peer Evaluation Rotation	73

Section D: Clinical Procedures

Competency Evaluation	75
Aseptic Technique/Infection Control	75
Doffing Checklist	77
Air Polishing	78
Caries Risk	80
Cleaning Removable Protheses	81
Dental Charting	81
Desensitizing	82
Disclosing Procedure	83
Ethics/Professionalism/Core Values	83
Extra/Intraoral Examination	84
Flossing	85
Fluoride Application	86
Implant Patients	86
Instrumentation	87
Instrument Sharpening	90
Intra Oral Camera Protocol	91
Medical and Dental History	92
Nutritional Counseling	93
Patient Education	93
Periodontal Charting	94
Phase-Contrast Microscope	94
Pit and Fissure Sealant Application	96
Polish/ De-plaque and Floss	97
Polishing – Seating Positions	98
Scaling and Root Planing	98
Tobacco Cessation Counseling	99
Oral Hygiene Instructions: Tooth Brushing	101
Treatment Plan/ Refusal of Treatment	102
Ultrasonic Scalers	102
Vital Signs	103

SECTION A

STUDENT INFORMATION

ADMINISTRATION

John A. Valenza, DDS	Dean and William N. Finnegan III Distinguished Teaching Professor in the Dental Sciences
Robert D. Spears, MS, PhD	Associate Dean for Student & Academic Affairs, Professor
Shalizeh Patel, DDS, MEd	Associate Dean for Patient Care, Professor
Muhammad F. Walji, MS, PhD	Associate Dean for Technology Services & Informatics, Professor
Lisa Cain, PhD	Associate Dean for Professional Development & Faculty Affairs, Professor
Arthur H. Jeske, DMD, PhD	Associate Dean for Strategic Planning, Professor
Mary.C."Cindy" Farach-Carson PhD	Associate Dean for Research, Professor
Joe Morrow, BBA	Associate Dean for Management
Nikola Angelov, DDS, MS, PhD	Chair, Department of Periodontics & Dental Hygiene, Professor
Victoria Patrounova, RDH, MHA	Director, Dental Hygiene Program, Associate Professor

UTHealth School of Dentistry Vision and Mission:

Vision

"Improving Oral Health ... Improving Overall Health"

Mission

"To improve human health by providing high-quality education, patient care, service and research in oral health for Texas, the nation and the world."

GENERAL GUIDELINES

UTHealth uses email as its primary and official method of communication. This is the policy of the Dental Hygiene Program and is in the best interest of the student. It is your responsibility to monitor your official UTHealth email on a daily basis.

A change in your name, address or telephone number should be reported promptly to the coordinator of the Dental Hygiene Program. Students are responsible for ensuring that their current address and telephone number are correct on myUTH. Updates to address information may be made on [myUTH](#). The [address change form](#) is also available at the Registrar's Office and Student Affairs Offices.

Permission of the Director and Office of Student Affairs is to be obtained before soliciting funds or conducting any type of campaign in the school. For more information, refer to HOOP Policy 165, Solicitation on Campus at <https://www.uth.edu/hoop/policy.htm?id=1448178>.

Smoking is not permitted on UTHealth premises. Smoking in public while in scrubs is considered unprofessional.

Dental Hygiene students are permitted in the clinic and laboratory only for officially scheduled activities or when such facilities are not being used by authorized groups.

The school is unable to provide secretarial services for students.

Students are expected to clear seminar rooms and cubicles by 4:50 p.m. each day. Students are not permitted in these areas on weekends or holidays.

A dental bookstore store offering supplies is located on the 2nd floor to allow students to purchase the necessary instruments and supplies as specified in the Student Instrument List. Textbooks are also available for purchase.

NOTE: Rules and regulations of The University of Texas Health Science Center at Houston and The University of Texas School of Dentistry at Houston apply to Dental Hygiene students. **Dental Hygiene handbook policies and guidelines are subject to change.**

DENTAL HYGIENE COMMUNITY LIAISON COUNCIL

The Dental Hygiene Program has a good relationship with the Houston area dentists and dental hygienists. A formal active liaison exists among these professional groups by way of the Dental Hygiene Community Liaison Council. Members of the Council meet annually to discuss current trends in dental and dental hygiene practice; and to assist in determining community health and dental hygiene employment needs. Health professionals, a second-year dental hygiene student, and the dental hygiene faculty serve on the council.

DISABILITY ACCOMMODATION

UTHealth ensures equal educational opportunity for all disabled individuals who are otherwise qualified, with or without reasonable accommodation.

If any student has questions about a disability or accommodation, or feels that he or she has been discriminated against on the basis of a disability, he or she should contact the UTHealth Office of Equal Opportunity and Diversity. Policies and procedures of UTHealth can be found in the UTHealth Handbook of Operating Procedures (HOOP). HOOP Policy 101, Disability Accommodations, can be found online at <https://www.uth.edu/hoop/policy.htm?id=1448050>

If you believe you have a disability requiring an accommodation, please contact:

Dr. Robert Spears
Associate Dean for Student & Academic Affairs
504 Disability Coordinator for UTSD
713-486-4166

For any questions or additional information, please call 713-500-2255 or email CALL@uth.tmc.edu.

Required Notice of Opportunity and Procedure to File Complaints with The Commission on Dental Accreditation (CODA)

The intent of this message is to inform students, faculty, constituent dental societies, state boards of dentistry and other interested parties that an appropriate, signed complaint (see definition below) may be submitted to the Commission on Dental Accreditation regarding any Commission accredited dental, allied dental or advanced education program.

Definition of Complaint:

A complaint is defined by CODA as one alleging that a Commission accredited educational program may not be in substantial compliance with Commission standards or required accreditation procedures.

- ❖ These issues and concerns may be discussed with the Associate Dean for Academic Affairs, at any time.
- |
- ❖ The Commission on Dental Accreditation will review complaints that relate to a program's compliance with the accreditation standards. The Commission is interested in the sustained quality and continued improvement of dental and dental-related education programs but does not intervene on behalf of individuals or act as a court of appeal for individuals in matters of admission, appointment, promotion or dismissal of faculty, staff or students.
- ❖ A copy of the appropriate accreditation standards and/or the Commission's policy and procedure for submissions of complaints may be obtained by contacting the Commission at 211 East Chicago Avenue, Chicago, Illinois 60611-2678 or by calling 1-800-621-8099, extension 4653.

I have read and I understand that I have the right to contact the Commission on Dental Accreditation if I so desire.

Print name _____

Signature _____

Date _____

August 20, 2018
Fall Semester

ADVISORY SYSTEM

The Dental Hygiene Advisory System allows development of student/faculty relationships and provides a faculty counselor that the student can contact for assistance. Each advisor is assigned a group of students at random from each dental hygiene class. The advisory groups remain fixed for the entire year. When changes are necessary, it is then the prerogative of the Director to transfer the student to another advisor.

The purposes of the advisory system are:

1. To allow faculty to facilitate, monitor, and assist in the attainment of students' individual goals;
2. To allow students the opportunity to discuss academic and clinical progress with a faculty member;
3. To encourage students to direct questions to a faculty member or share problems that may affect scholastic performance; the advisor may refer the student to outside agencies for additional assistance; and
4. To allow the advisor to monitor the students' progression or regressions in their professional development and to plan remedial or enrichment work accordingly.

PROTOCOL FOR RESOLUTION OF STUDENT CONCERNS

In general, student complaints are governed by Hoop Policy 220, Student Complaints, and more information on this policy can be found at <https://www.uth.edu/hoop/policy.htm?id=2553c1c1-c490-4ad0-a570-e263e12e0dff>.

In the event a student has a concern with a dental hygiene faculty member, the following protocol must be followed:

- The student must make an appointment with the faculty member during their posted office hours and discuss their concern/s;
- If after the appointment with the faculty the student feels the concern has not been adequately addressed, the student must email the dental hygiene program director requesting her/him to meet with the faculty member about the concern/s;
- After the dental hygiene program director meets with the faculty, she/he will provide the student an opportunity to meet with both the faculty and herself to discuss and resolve the concern/s.

In the event a student goes to a faculty member about a concern with another faculty member, faculty will not discuss the concern with the student but rather refer the student to this required program protocol.

DENTAL HYGIENE COMPLAINT, SUGGESTIONS, & COMPLIMENTS FORM

Students can also express their concerns, complaints, compliments, and suggestions with total anonymity. Access the form at the following link:

https://uthmc.az1.qualtrics.com/jfe/form/SV_0DjFupYHtCkYGB8

The dental hygiene program director meets with the faculty or student to discuss and resolve the concerns. Summary of discussion will be submitted to a secure file.

ETHICS

Ethics is the part of philosophy that deals with moral conduct and judgment. There are several principles that health care professionals must be aware of in the practice of their profession. The major principles are:

- *Autonomy* – self-determination in a person; the right to participate in and decide on a course of action without undue influence
- *Beneficence* – promoting good or well-being

- *Non-maleficence* – the duty to avoid harming the patient, summarized by the phrase “do no harm”
- *Confidentiality* – the precept by which information shared by a patient during the course of receiving health care is kept in confidence by the health care provider
- *Veracity* – a duty to tell the truth when information is disclosed to patients about treatment
- *Societal Trust* – maintaining a bond of trust in the relationships between healthcare professionals, patients, and the public`
- Professionalism is defined as “the conduct, aims, or qualities that characterize or mark a profession or a professional person”. While these characteristics may vary from profession to profession, the practice of dental hygiene requires professionalism which goes far beyond basic honesty and integrity.

While you will be evaluated on the basis of your ethical behavior, professionalism and intellectual and psychomotor abilities, you are also required to be exemplary in your grooming, personal conduct, and relationships with faculty, peers, and patients.

The students’ responsibilities at UTHHealth and UTSD may be classified in five broad areas, which are as follows:

- Academic performance
- Academic integrity
- Professional conduct
- Conduct associated with the university, but not directly related to academic or professional training of the student
- Off-campus conduct, not associated with the university, which may reflect adversely on the image and reputation of the University, including the student’s social media activity.

The faculty and administration are dedicated to the development of professionalism in all School of Dentistry students. The aim of the institution is to create a learning environment which offers students the opportunity to develop standards of excellence which will sustain them throughout their professional careers. For more information, refer to Hoop 186, Student Conduct and Discipline at <https://www.uth.edu/hoop/policy.htm?id=1448220>.

DENTAL HYGIENE LICENSURE ELIGIBILITY

According to Laws of the State of Texas and the Texas State Board of Dental Examiners a person applying for initial licensure to practice Dental Hygiene in the State of Texas may be ineligible for licensure due to a previous conviction or deferred adjudication for a felony or misdemeanor offense. Reference rule 101.8 at the following link:

[https://texreg.sos.state.tx.us/public/readtac\\$ext.TacPage?sl=R&app=9&p_dir=&p_rloc=&p_tloc=&p_ploc=&pg=1&p_tac=&ti=22&pt=5&ch=101&rl=8](https://texreg.sos.state.tx.us/public/readtac$ext.TacPage?sl=R&app=9&p_dir=&p_rloc=&p_tloc=&p_ploc=&pg=1&p_tac=&ti=22&pt=5&ch=101&rl=8)

ACADEMIC INTEGRITY

It is imperative that students maintain high standards of integrity in their scholastic endeavors. It is the responsibility of the faculty to see that such standards are maintained. Scholastic dishonesty is the submission, as one’s own, of material that is not one’s own. As a general rule, it involves, but is not limited to, one of the following acts: cheating, plagiarism, and collusion.

Cheating is defined as receiving unauthorized aid on an examination, quiz, paper, clinic or laboratory experience or course project such as:

- Copying from another student's test paper or laboratory project.
- Using unauthorized materials such as class notes, crib notes, etc., during any examination or quiz. The presence of textbooks and/or other course material such as class notes, crib notes, etc. is prohibited for that test unless specifically allowed by the course instructor.

- Knowingly using, buying, stealing, transporting, soliciting, disseminating or attempting to disseminate the contents of an unreleased exam.
- Unauthorized collaboration or attempting to obtain unauthorized aid from another student during a test.
- Unauthorized substitution, or allowing another to be substituted for oneself, to take or perform a test or a laboratory procedure, or to sign class attendance records for another, or to disseminate access passwords for the quiz or exam.
- Giving or accepting money or other inducements to obtain an unreleased test or information about an unreleased test.
- Unauthorized collaboration on a take-home assignment that is not specifically permitted by the instructor or course director.
- Unauthorized or improper use of any technological or communication tool during an exam.
- Performing unauthorized procedures outside of clinical hours or at an outside facility.
- Performing another student's preclinical /clinical projects.

Plagiarism is the appropriating or obtaining by any means of another's work, and the unacknowledged submission or incorporation of it into one's own written work then offered for credit. Plagiarism includes, but is not limited to:

- Failing to give credit for ideas, statements of facts or conclusions derived by another author; failure to use quotation marks when quoting directly from another, whether it is a paragraph, a sentence, or part thereof; failure to properly cite the work of another person.
- Submitting a paper purchased or obtained from a "research" or term paper service or individual supplier, or otherwise buying or receiving such work.
- Submitting a paper, in whole or in part, obtained from an internet resource.
- Giving a speech or oral presentation written by another and claiming it as one's own work.

Collusion means the unauthorized collaboration with another person in preparing academic assignments which are offered for credit.

Penalty for Scholastic Dishonesty

The penalties for scholastic dishonesty, as described in HOOP Policy 186 found online at <https://www.uth.edu/hoop/policy.htm?id=1448220>, may include, but are not limited to: disciplinary probation, withholding of transcript or degree, being barred from readmission; failing grades for the exam, practical, and/or course; denial of degree; repetition of the year; suspension from the institution; or expulsion from the institution.

Reporting a Suspected Breach

Any member of the UTHealth School of Dentistry community who has reasonable cause to believe that a breach of this Code of Academic Integrity has been committed, has an ethical obligation to:

- Report the incident via the "UTHealth School of Dentistry Academic Integrity Reporting Form" (see below)
- Inform the course director/instructor, the UTSD Associate Dean for Student and Academic Affairs, or The Office of Institutional Compliance of the suspicions and the reasonable basis for them. This also includes self-reporting. Anonymous reports can be made to The Office of Institutional Compliance via their website (<http://www.lighthouse-services.com/compliance.uth.tmc>) or by calling 1- 833-222-0056. Suspected breaches of academic integrity will be reported to the UTSD Associate Dean of Student and Academic Affairs.

Reports of suspected breaches of this Code will be addressed in accordance with the policy and procedure described in HOOP 186 and the School's *Clinic Manual*, *DDS Student Guide to Academic Affairs*, *Dental Hygiene Student Handbook*, and *Advanced Education Manual & Course Catalog* documents, as applicable.

UTHealth School of Dentistry at Houston Academic Integrity Reporting Form

Date: _____

Reporting Person (optional):

Name: _____
Last First MI Contact #: _____

Course Name / Number: _____

Student(s) Involved in Alleged Incident: _____

Date of Alleged Incident: _____

Please provide a detailed description of the alleged incident:

Report Received By (name): _____

Date Report Received: _____

PROFESSIONALISM

Students are expected to perform in a professional and ethical manner in all aspects of the delivery of patient care. The School of Dentistry responds to inappropriate clinic performance or behavior (“infractions”) by students through academic corrective action. Infractions which are deemed more serious in nature (“cardinal”) may be referred to the appropriate Evaluation & Promotion Committee or the Associate Dean for Student and Academic Affairs. Investigation and, if indicated, appropriate action taken will be in accordance with policies described in the DDS Student Guide to Academic Studies with UTHealth HOOP Policy 186 Student Conduct and Discipline. Students are responsible for knowing and observing state and federal laws as well as the policies of both UTHealth and the School of Dentistry. [Students can find the full listing of UTHealth Handbook of Operating Procedures online at https://www.uth.edu/hoop/index.htm.](https://www.uth.edu/hoop/index.htm)

Unprofessional behavior will not be tolerated. The first occurrence will result in a warning and deduction of points. Examples of unprofessional behavior include, but are not limited to:

- Audible signals emitting from devices or cellular phones during lectures, or other presentations
- Use of cellular phones or web devices for other than class activities during class
- Leaving class after the presenter has started or before the presenter has concluded
- Engaging in audible conversations with colleagues during class presentations
- Failure to adhere to the dress code as defined in the current *Dental Hygiene Program Handbook*
- Unprofessional behavior, attitude or language (rudeness, profanity, etc.)
- Intentional physical destruction of equipment or building facility

CORRECTIVE ACTION

The Associate Dean for Patient Care and Assistant Dean for Clinical Education are responsible for executing UTHealth/UTSD policies as they relate to the clinical academic program and patient care. As all clinical faculty share responsibility for student compliance of clinic policy and procedure, most minor clinical infractions may be resolved by an attending faculty member. However, in those instances where, in the judgment of a faculty or appropriate staff member, referral to clinic administration for assessment, investigation and possible corrective action is indicated, the following persons will be responsible for determining the appropriate actions as follows:

For infractions involving a dental hygiene student, the Director of the Dental Hygiene Program is to be contacted immediately. Further referral, if indicated, would be to the Dental Hygiene Student Evaluation & Promotion Committee or the Associate Dean for Student and Academic Affairs.

Faculty, students or staff may report infractions by completing a *Student Incident Report*, which is available on Canvas in clinical courses. Reports should be completed and returned as noted above.

STUDENT INCIDENT REPORT

Date of Incident: _____

Location: _____

Name of Student(s) Involved: _____

Name of Faculty or Staff Member(s) Reporting Incident: _____

Reason for Report:

- Attendance
- Patient treatment
- Policy/Procedure
- Professionalism
- Other: _____

Comment: _____

Student Signature: _____ Date: _____

Faculty or Staff Signature: _____ Date: _____

DISTRIBUTION:
WHITE – Office of Clinical

Education

(Return this form to Assistant Dean for Clinical Education, SOD 3510)

CLINICAL INFRACTIONS

Clinical infractions which warrant corrective action may include, but will not be limited to:

- Violations of an ethical and/or professional nature
- Failure to comply with infection control protocol (See UTSD Clinical Manual Section 3 – <https://inside.uth.edu/dentistry/clinical-resources/clinic-manual/#panel44f16b76-4a8a-4488-96cb-fb2912354228-3> for additional details regarding infection control infractions)
- Improper management of patient records
- Misuse or inappropriate use of electronic patient record, including downloading or unprotected printing of PHI
- Removal of patient-sensitive information from the School of Dentistry building
- Use of a password or identification card that is not the student's own
- Use of instruments or materials not approved for use in SD clinics
- Misuse of instruments or materials, or failure to return excess unused materials
- Failure to attend and successfully complete required annual clinical updates
- Failure to attend mandatory clinical meetings, exercises, or assignments

Violation of clinic policies will be dealt with on an individual basis. In general, and dependent upon the severity of the infraction, violations will carry one or more academic penalties unless otherwise stated, and not necessarily in the following order:

- Students are allowed 2 warning incident reports. Required action by the student receiving 2 clinical incident reports may include writing of a report and/or presentation related to the infractions.
- The 3rd incident report goes into the student's academic record and the student is suspended from clinical and rotation for a minimum of one (1) week and for as long as necessary for remediation in the area of the infraction.
- Students on clinical suspension may continue to access the EHR and view patient records. They may **not**, however, perform the following: schedule, treat, assist or observe patients in clinic and cannot obtain clinical instruments, equipment or materials from clinical dispensaries.
- If a student suspension led to failure of a clinical course, a decision on the dismissal of the student goes through the E&P Committee.

CARDINAL INFRACTIONS

Cardinal Infractions are considered serious in nature. Examples of cardinal infractions include:

- Providing treatment outside the Dental Hygiene Scope of Practice as determined by the Texas State Board of Dental Examiners.
- Any infection control infraction which seriously endangers the health of a patient, student, faculty, or staff.
- Using non-sterile handpieces, instruments or other items, when sterility is required.
- Disposing of any red biohazard bag inappropriately.
- Handpiece or instruments not returned for sterilization within 24 hours.
- Verbal or physical misconduct involving a patient, student, faculty, or staff member
- Falsification of a patient record or clinic document
- Abandonment of a patient

Students face automatic and immediate suspension from the clinic, if deemed appropriate, until such reasonable time as a final course of action is determined by the Dental Hygiene Student Evaluation & Promotion Committee.

Documentation of violations of clinic policy will be maintained in appropriate University files.

ATTENDANCE

Attendance is expected at all scheduled lectures, clinics, laboratories, seminars, case presentations, rotations, and individual faculty appointments. Attendance will be taken at each lecture/lab. As a matter of courtesy to the speaker and to get maximum benefit from the lecture/lab, you should make every reasonable effort to arrive at class before the presenter begins the lecture. Attendance is considered one measure of a student's **professional conduct**. Students who abuse attendance requirements will be considered for academic action. All excused absences must be approved by the Director of the Dental Hygiene Program. To receive an approved excused absence the student must submit appropriate documentation to the Dental Hygiene Office **within 3 days** upon return to class/clinic. This will allow the student to make up missed exams.

At the discretion of the course director, attendance may be taken through the use of various methods such as sign-in rosters, audience response apps and/or quizzes. Tardiness or leaving class early may be counted as an absence. Attendance records are official school documents, and thus falsification of these records by any student will constitute a significant act of dishonesty. At the discretion of the course director, attendance may play a part in the course grade as described in the course syllabus.

RELIGIOUS HOLY DAYS

“Religious holy day” means a holy day observed by a religion whose places of worship are exempt from property taxation under Section 11.20 of the Tax Code.

The University of Texas School of Dentistry at Houston, in compliance with Section 51.911 of the Texas Education Code, will allow a student who is absent from classes for the observance of a religious holy day to take an examination or complete an assignment scheduled for that day within a reasonable time after the absence if, no later than the fifteenth day of the semester, the student notifies the course director of each class the student has scheduled on that date, that the student will be absent for a religious holy day. Students are responsible for notifying, in writing, the course director of each class. A copy of the letter must also be provided to the Office of Student and Academic Affairs. The course director will establish a reasonable date for the completion of the assignment or examination and notify the student prior to the aforementioned holiday.

SEVERE WEATHER AND SCHOOL CLOSURE

UTHealth has a resource to provide emergency alerts in instances of severe weather, university status change, or if there's an imminent threat to the community. Participants can opt in to UTHealth Alert by providing their phone number to the database. They can change their number or opt out of the service at any time. Instructions on how to enroll or update your cell phone number can be found on the UTHealth Alert website at <https://www.uth.edu/uthealthalert/enroll>.

In the case of an unanticipated absence necessitating cancellation of patient appointments, it is the student's responsibility to notify their patients.

In case of emergencies contact the UT Police Department by dialing 911; identify yourself as a student of the University of Texas School of Dentistry and give them your location (identify the streets that intersect closest to your location). For non-emergencies contact UT Police at 713-792-2890.

PROCEDURES FOR REPORTING ABSENCES

Students are responsible for informing the educational coordinator of the Dental Hygiene Program at 713-486-4084 and course directors if unable to attend school all day or part of it. If it is necessary to leave school early for the day, the educational coordinator must be notified. Students are responsible for contacting course directors regarding assignments prior to an absence (if known) or after the absence.

In the case of an unanticipated absence necessitating cancellation of patients, it is the student's responsibility to notify the patients, clinic coordinator, clinic course director and the department coordinator by 8:30 a.m. on the day of the absence. Absences reaching **three or more days** require a physician's letter or other suitable documentation for the absence.

It is the student's responsibility to contact the course directors of missed classes **within three days** of return to the school to determine what, if any, arrangements are to be made for missed coursework (examinations, practical exams, etc.). If a scheduled examination, quiz, or required activity will be missed, the course director should also be contacted, preferably before the scheduled start of the examination or required activity.

Anticipated absences, e.g. interviews, doctor appointments, etc., should be discussed with the appropriate course directors prior to the absence so that arrangements can be made as needed.

UTHEALTH ID BADGES

UTHealth ID badges are **required** to be visibly worn at all times by students, staff, and faculty when in the UTHealth area buildings. Individuals who are not wearing valid ID badges or are unable to produce them upon request may be asked to leave the building. ID badges are used for entrance into the building, and are used to check out books from the UTSD Library. The replacement fee for a lost or damaged identification badge is \$10.00.

APPEARANCE GUIDELINES

Each individual involved in the Dental Hygiene Program reflects on how others view the program, the School of Dentistry and the profession. The attitude, mannerism, and physical bearing displayed in relationships with patients, staff, faculty, classmates and the public are a serious responsibility which we must exhibit in an ethical, safe, and professional demeanor. Furthermore, patients often form a first impression based on the physical appearance of their healthcare provider and develop trust more quickly with a person who portrays cleanliness, and professionalism. The following guidelines are given to assist all persons in understanding and accepting this responsibility.

Students, faculty, staff, and administration are all responsible for assisting one another in being good influences on the perception of our program to each other, School of Dentistry administration and the community. These guidelines are not presented to cover every detail and situation. It is expected that individuals know proper and appropriate behavior, and that they will support it by their own example.

A. Clinic/Classroom Attire

Clinic Attire— Appropriately colored scrubs (as required by the School of Dentistry) are the only attire permitted. Scrub suit colors differ for each class and will remain the same color until the student graduates. Scrubs are to be in good repair, neat in appearance, free of stains, wrinkle-free with the top and pants colors matching. Surgical scrubs must NOT be worn with street clothes. However, t-shirts may be worn as an undergarment with scrub tops. For personal comfort outside of the clinic, matching scrub jackets or professional white coats may be worn over surgical scrubs.

A clinic gown must be worn over scrubs while in the clinic, radiology, x-ray processing room and patient assessment area. A clinic gown must be changed after treatment of each patient (or sooner if visibly soiled). Clinic gowns must not be worn outside patient treatment areas.

Shoes: Appropriate clinical shoes are to be worn in the clinical environment at all times. Shoes must be clean and in good repair. Shoes must be made of a wipe-able material, typically leather-like material is most appropriate. Shoes must be closed toe and have a back. Shoes with holes on the top, known as “Crocs” or soft fabric shoes such as “Toms” may NOT be worn in the clinics for safety and infection control reasons.

Socks: OSHA principles direct that skin is to be covered if there is the likelihood of exposure to chemicals and/or bodily fluids. Therefore, socks must cover the ankle area (no skin visible when seated) and are included as a part of the complete uniform. Socks should be white, black or matching the color of the scrubs. However, patterned/contrasting color socks are acceptable as long as the pattern is not considered offensive (e.g., no crude or vulgar language is imprinted on the socks).

Skirts: Female students are allowed to wear appropriate, class-colored clinical scrub skirts for religious purposes. The student’s legs must be covered while in the clinic setting. If the skirt is not floor length, the student’s legs must be covered with opaque tights or leggings. Tights or leggings should be black or matching the scrubs.

Hats, Scarves and Religious Headwear: Hats, sports bands, and headbands are not allowed as they could collect microorganisms and create the opportunity for cross-contamination. Religious accommodations for head coverings may be requested by following the guidance provided in [HOOP 112](#). Head covering needs to be secured in a manner which will not jeopardize student or patient safety, and these items must be washed after each use.

A disposable or washable head covering that meets infection control guidelines will be worn in any situation where there is the probability of exposure to bodily fluids as occurs with splash, splatter, or aerosols such as ultrasonics and air polishing.

Hair: should be kept clean and well-groomed. While there are no restrictions on hair length, hair must be pulled back and kept away from the treatment field, so it does not require handling during any clinical procedure. Facial

hair must be clean, neatly trimmed, and well-groomed to fit under the surgical mask to prevent contamination. When wearing an N95 respirator, facial hair must be cleanly shaved and individuals must be fit-tested and trained to properly wear the N95 device and ensure protection is provided. Students who have facial hair or head coverings for religious purposes must communicate this exception to the program director prior to clinical activities.

Eyewear: Protective glasses with side shields and surgical masks with a face shield and surgical mask will be worn in all clinical situations which expose the clinician to aerosol and splatter. Therefore, artificial, long eyelashes are discouraged as they may pose interference with protective eyewear and/or loupes.

Jewelry: Rings (except smooth surface wedding rings), bracelets, long necklaces, and long dangling earrings are not allowed while treating patients. Jewelry harbors microbes, interferes with the ability to perform successful hand hygiene, compromises the integrity of gloves, and can serve as a fomite to transmit infection beyond the operatory. Earrings should not extend beyond the ear. Facial piercings are limited to small nose piercings that must be covered with a mask while in clinic. Gauge, plug earrings or larger than normal holes in the ears will be covered with a disposable bouffant cap when in the clinic. Wristwatches are allowed when covered by the clinic gown.

Tattoos: They must be covered if they are visible on the face/neck and/or have content that is inappropriate (contains obscene/offensive images, designs, or language). Fresh tattoos must be covered with a bandage to reduce risk of infection while in the lab or clinical setting.

UTHealth Identification Badges: Students are required to have the UTHealth I.D. badges at all times. When in the clinic, your badge must be placed in a pocket under the gown.

Writing pens are not to be attached to the outside of the clinic gown.

Gum chewing is **NOT** acceptable in clinic or other professional settings. Chewing gum in the clinical setting is unsanitary and unprofessional. ***INFRACTIONS OF THE DRESS CODE WILL BE REFLECTED IN THE PROFESSIONALISM DAILY GRADE.***

Classroom – Besides full scrubs, only **UTHealth t-shirts** with scrub bottoms are considered appropriate attire in classes or preclinical laboratories. Being allowed to wear UTHealth t-shirts in class is a privilege and not a right. If any student abuses this privilege by wearing a non-UTHealth t-shirt, this privilege will be taken away from ALL students. Students may wear their white School of Dentistry white coat or a sweater/fleece jacket over scrub tops in the classroom.

Other times- On infrequent occasions AND upon approval from the program director, appropriate business attire may be worn (dress pants and a dress shirt or blouse). Jeans, shorts, leggings, t-shirts, tank tops, workout clothing, garments with hoods, or other non-professional attire are not allowed at school or school-related functions. These personal appearance standards are in effect from 7:00 am to 6:00 pm Monday through Friday. Students who are not in compliance with these minimal dress requirements will not be allowed to remain in the building.

B. Personal Hygiene

Close proximity with patients requires meticulous personal hygiene at all times. It is necessary to bathe daily and use a dependable deodorant/antiperspirant. Strong perfumes or colognes are to be avoided.

Fingernails must be kept short enough (less than ¼ inch long) to allow thorough cleanliness. The fingernails should not extend beyond the fingertips to prevent collections of microbes and tears in gloves and to allow easy

hand cleaning. Nail polish is allowed however it must be kept fresh and unchipped, as chipped nail polish has been shown to harbor and increase the number of bacteria.

False fingernails, including acrylics, overlays, tips, and shellac or “gel” nails are prohibited. (CDC Guidelines <https://www.cdc.gov/hygiene/personal-hygiene/nails.html>)

Plaque control must be practiced; mouth odors must be controlled. If you have a problem with your breath, brush your teeth and tongue and use a breath spray, mouthwash, or breath mint prior to seating your patient.

EXAMINATIONS

Examinations are administered by the course directors or their designee. The date, time, and location of major exams are published in the syllabi and the student schedule. If a student does not attend an examination on the specified date and time, the student must contact the course director within five working days of the exam date. If the student cannot produce acceptable documentation for an excused absence to the Office of Student and Academic Affairs, a grade of zero (“0”) will be recorded. A student with an unexcused absence may be granted a make-up exam; however, a penalty will be applied.

Students producing adequate documentation to the Dental Hygiene Program Director and the Office of Student and Academic Affairs will be granted an “excused absence” and allowed to take a make-up exam, without penalty, to be administered by the course director. The format of the make-up exam is at the course director’s discretion and it may be different than the original exam. The course director, or their designee, will grade the exam.

All exams are required to be graded within seven days following the examination. Grades will be posted in Canvas

1. No student is permitted to leave the examination room before completing the examination, unless the instructor’s permission has been granted.
2. No student is permitted to enter the examination room to begin an examination after another student has completed the examination and left the room.
3. Computer generated examinations may be given online or on computers in testing rooms or labs.
4. All books, purses, and electronic devices must be left at the front of the room during an examination or outside the examination room. Students will not be allowed to have any of these materials at their desks during the exam.
5. If you are/were absent when a major examination will be or was given, it is your responsibility to notify the course director.
6. Make-up examinations may be rescheduled after finals have been given as determined by the course director.
7. Students may be recorded while taking exams within the School of Dentistry building or when taking an exam off campus.
8. If a student is found cheating, disciplinary action will be taken.
9. Approved security screens are required for all graded exams and quizzes taken electronically

ACADEMIC STANDARDS

GRADING SYSTEM FOR CLINIC AND CLINIC RELATED COURSES:

93 – 100 = A
84 – 92 = B
75 – 83 = C
<75 = F

GRADING SYSTEM FOR NON-CLINIC RELATED COURSES:

90 – 100 = A
80 – 89 = B
79 – 70 = C
<70 = F

Passing

Grades for didactic and clinical courses are letter grades. A minimum grade of C will be required in all courses and an overall average of C (2.00 GPA) must be maintained. Student Organization officers must maintain a 2.75 GPA to remain in office. **An acceptable level of clinical proficiency must be demonstrated in each clinic before the student will be permitted to begin the next clinic.**

Failing

A grade of 69 or below designates failing work in non-clinical courses; a grade of 74 or below designates failing work in clinic-related courses.

Incomplete

A grade of incomplete (I) may be given under rare circumstances and only upon approval by the Director of the Dental Hygiene Program and the Evaluation and Promotion Committee. A grade of incomplete may be either incomplete while passing or incomplete yet failing. A grade of Incomplete yet failing generally results in a Final course grade of F.

GRADE GRIEVANCE PROCEDURE

In attempting to resolve any student grievance regarding grades or evaluations, it is the student's responsibility to arrange an appointment to discuss the grievance with the faculty member /the appropriate course director within five working days of the grade in question. Individual faculty members retain primary responsibility for assigning grades and evaluations. A faculty member's judgment in such cases is final unless there is substantial evidence of discrimination, differential treatment, or error. If, after meeting with the appropriate faculty member, the student feels that the grade grievance has not been adequately addressed, the student may appeal the grievance in writing to the Associate Dean for Student and Academic Affairs within seven working days. The Associate Dean for Student and Academic Affairs will review the case and submit a written recommendation to the Dean within fourteen working days. The Dean will respond in writing to the student's grievance within five working days. In academic issues the determination of the Dean is final and not subject to further appeal.

WARNING, PROBATION, AND DISMISSAL

Warning: Students will receive a letter of warning at mid-semester for unsatisfactory progress in didactic, laboratory, or clinical courses. Students will be expected to show sufficient improvement with a passing grade in those areas of deficiency by the end of that semester to avoid being placed on probation or considered for dismissal. In addition, the student is expected to satisfactorily progress in the other courses in the curriculum.

Probation: Students having a semester GPA or cumulative GPA below 2.00 will be placed on probation if not dismissed from the program. Students who have been placed on probation must show acceptable improvement and satisfy the conditions of the letter placing them on probation within the following semester, or they may be dismissed for academic reasons. Students placed on probation may be ineligible for financial aid and will be ineligible to hold class or SCADHA offices. Reinstatement is at the discretion of the Dental Hygiene Program Director.

Dismissal: Students will be considered for academic dismissal if they have a cumulative grade point average below 2.00 at the end of a semester. Students will be considered for academic action that could include dismissal if they have one or more failing course grades in a given semester or in more than one semester.

If a student demonstrates the inability to progress either didactically or clinically, he/she will be considered for dismissal from the Dental Hygiene Program by the Evaluation and Promotion Committee – Dental Hygiene Subcommittee. The decision will be made by the committee members at a meeting held at the end of the semester. Specific guidelines for academic dismissal are listed above.

ACADEMIC ACTION AND APPEAL PROCESS

A University of Texas School of Dentistry student may appeal any academic action by the Evaluation and Promotion (“E & P”) Dental Hygiene subcommittee to the Associate Dean for Student and Academic Affairs, in writing, within three calendar days after receipt of notice of the academic action. The student must provide the Associate Dean for Student and Academic Affairs a “complete” appeal, which includes at least a written statement clearly explaining all rationale for the appeal and any additional documentation the student possesses that the student believes supports the student’s rationale for the appeal.

The Associate Dean for Student and Academic Affairs will refer each complete appeal to an Ad Hoc Appeal Committee (“Appeal Committee”). The Office of the Associate Dean for Student and Academic Affairs will assist by scheduling the meetings of the Appeal Committee.

- The Chair of the Appeal Committee will be selected and appointed by the School of Dentistry Committee on Committees and approved by the Faculty Senate (an alternate Chair will also be selected from among the faculty of the School of Dentistry). The Chair will preside over the Appeal Committee. The length of the Chair’s term will be three years. The alternate will preside over the Appeal Committee in the event that the Chair is unable to attend.
- The Appeal Committee will be made up of the chairs of each of the E & P subcommittees not involved in the academic action being appealed. Vice chairs of the E & P subcommittees may serve in this role in the event a subcommittee Chair is unable to participate. In addition, an additional member of the Appeal Committee will be selected by the Associate Dean of Academic Affairs from among SOD faculty. This member of the Appeal Committee cannot be the student’s faculty advisor or a member of the E & P subcommittee making the decision being appealed.
- Each of the Appeal Committee members will have one vote. In the case of a tie vote, the Chair of the Appeal Committee will vote to break the tie.

The Appeal Committee will review the student’s appeal letter and/or written statement and documentation, if any, submitted by the student, meet with the student, the student’s faculty advisor, the Chair of the E & P subcommittee taking the academic action being appealed, and other individuals at the discretion of the Chair of the Appeal Committee. The Chair of the Appeal Committee shall submit a final recommendation to the Dean within seven calendar days of the final Appeal Committee meeting. The Dean shall consider the recommendation of the Appeal Committee, may review the materials submitted to the Appeal Committee, and may interview other individuals. At his or her discretion, the Dean may meet with the student. The student will be notified of the Dean’s decision within 10 calendar days after the Dean’s receipt of the Appeal Committee recommendation. The Dean’s decision regarding the academic action of the E & P subcommittee is final. The results of the Appeal Committee may be shared with the Chair of the Department involved.

The student, upon written request to and approval in writing from the Associate Dean for Student and Academic Affairs, may continue academic studies while the appeal of an academic action is under review and until the student receives notification of a final decision by the Dean.

If after the appeals process is completed an academic action of dismissal is upheld, a dismissed student must immediately discontinue participating in all UTSD educational activities. All personal belongings must be removed from the School of Dentistry facilities immediately upon receipt of the final decision of the Dean. If the decision is to repeat the year then the student must arrange for enrollment, financial payments, registration, and the removal of any holds on their records.

The School of Dentistry Student Evaluation and Promotion Committee consist of four subcommittees: The First Year Dental Student Evaluation and Promotion Subcommittee, the Second Year Dental Student Evaluation and Promotion Subcommittee, the Third/Fourth Year Dental Student Evaluation and Promotion Subcommittee, and the Dental Hygiene Student Evaluation and Promotion Subcommittee. Each subcommittee is led by a chair and a vice chair.

REMEDIATION

Students who fail exams or competencies will have the opportunity for remediation to assist them in improving their knowledge, skills and understanding during the regular semester. Students who have remediated a failed course, will still be required to pass the course in a similar manner stated in the course syllabus.

THE UNIVERSITY OF TEXAS HEALTH SCIENCE CENTER AT HOUSTON OFFICE OF STUDENT FINANCIAL SERVICES POLICY REGARDING SATISFACTORY ACADEMIC PROGRESS (SAP)

SAP will be reviewed annually after the end of each spring term and after all grades have been posted. SAP for entering students will not be reviewed until after completion of at least one semester.

- A student must maintain a cumulative GPA of 2.0 or greater at the end of each academic year. Those below the standard cumulative GPA will not be eligible for financial aid

Student Financial Services Policy on Satisfactory Academic Progress can be found online at the following link:

<https://www.uth.edu/dotAsset/67612ff0-aa4f-4e2c-b2d6-19653eafe106.pdf>

STUDENT WITHDRAWAL

Any UTSD student who does not intend to continue as a student must officially **withdraw**, rather than simply stop attending classes and laboratories. Students who decide to withdraw must complete a Checkout Sheet and a Student Exit Form, which are available in the Office of Student & Academic Affairs (Suite 4120). Following an exit interview, the student must secure clearance from the various offices noted on the Checkout Sheet and return it to the Office of Student & Academic Affairs. Completion of this process constitutes an official withdrawal.

LEAVE OF ABSENCE

Any student in good academic standing who needs or wishes to stop attending classes and laboratories temporarily, intending to continue studies at a later date, must submit a written request for a **leave of absence** to the Director of the Dental Hygiene Program and then the Associate Dean for Student and Academic Affairs stating the reason for the request, the length of leave requested, and the date for resuming studies. The request for leave of absence must be presented within five days of the attended start date of the leave.

The Associate Dean for Student and Academic Affairs will confer, when necessary, with the Associate Dean for Clinical Education regarding the leave request. The Associate Dean for Student and Academic Affairs will review the leave request and the student's academic record, and will recommend whether the leave should be granted and any conditions, which must be met for the student to re-enroll. In compliance with UTHHealth policy and procedures, additional documentation may be required for students seeking medical leave or for those students called to active duty military service.

A Leave of Absence will not be granted solely because of poor academic performance. For students in academic jeopardy, the Associate Dean for Student and Academic Affairs will refer the leave request to the Evaluation and Promotion Committee. The Student Evaluation and Promotion Committee will recommend to the Associate Dean for Student and Academic Affairs whether the leave should be granted and, if so, the point in the curriculum where the student may re-enter and any necessary remediation activities following re-enrollment.

The Associate Dean for Student and Academic Affairs will notify the student, in writing, of the action on the student's request, including any conditions, which must be met by the student, and the expected re-entry date. Following approval by the Associate Dean for Student and Academic Affairs, the student must complete a Checkout Sheet and a Student Exit Form, which are available in the Office of Student & Academic Affairs (Room 4120). The student must secure clearance from the

various offices noted on the Checkout Sheet and return it to the Office of Student and Academic Affairs. Completion of this process constitutes an official leave of absence.

In general, students can request a LOA for no longer than one-year, and those taking a leave of absence will re-enter the curriculum no later than the point at which the leave began, and students may be required to repeat a portion of the curriculum, or potentially the entire academic year for extended leave situations of less than one academic year. Students on leave from The University of Texas School of Dentistry at Houston for more than one calendar year may be required to repeat all or a significant portion of the curriculum.

A student on LOA must inform the Associate Dean for Student and Academic Affairs in writing of their intent to re-matriculate at least two calendar months prior to the anticipated re-matriculation date. Notification must include any supporting documentation required by conditions of the leave. Notification of a student's intent to return does not guarantee a right to return. Readmission is at the sole discretion of the appropriate E&P committee. A student who fails to return within the allotted 12 months of the leave or fails to notify the school will be administratively withdrawn from the program. Students on an approved LOA are not required to register for courses and are not required to pay instructional or other use fees. Access to UTHealth facilities and services may be restricted.

READMISSION

A student who voluntarily withdraws or is dismissed from the dental hygiene program in good standing and subsequently reapplies for admission will be considered on an individual basis by the Dental Hygiene Admissions Committee.

Requirements that govern the readmission of applicants to the dental hygiene program are as follows:

- The student must not have been out of dental hygiene school for more than five years at the time of acceptance.
- Readmitted students will be required to audit previously taken courses if more than three years or if major course revisions have occurred since their enrollment and must complete all course requirements satisfactorily.
- An interview will be required prior to an offer of admission.

ACADEMIC COUNSELING

Advanced Academic Training (AAT) is designed to help entering first-year healthcare students master their rigorous academic programs. Students are encouraged to call the UTHealth School of Dentistry, Office Educational Research and Development at 713-4125161 or email Dr. Ronald Johnson at Ronald.Johnson@uth.tmc.edu for individualized help in all aspects of their educational training.

CLINICAL EQUIPMENT INFORMATION

1. Students are responsible for the cleanliness of lockers, laboratory benches, and the laboratory in general, and clinical cubicles to which they are assigned. Physical plant personnel empty waste baskets each evening.
2. Turn off laboratory and cubicle lights, gas, water, and air when not in use. Place operator's stool in its original location.
3. Students are to inspect all equipment that may be missing or not in good working order at the beginning of each clinic session. At any time when equipment is missing, damaged, or malfunctioning, it should be reported to your instructor and to Educational Support Services, Room 3450, Telephone 713-486-4441. Computer problems should be reported to the HELP Desk at 713-500-4848. **DO NOT** use the item until it has been repaired.
4. Students are responsible for all equipment loaned to them (Cavitron, instruments, Piezo handpiece, ultrasonic inserts/tips, curing light, slow speed handpiece, etc.).
5. Any damage or loss will result in payment by the student to repair or replace loan items.

INSTRUMENT RETURN/ REPLACEMENT POLICY

In the event dental hygiene instruments are found to be dull, broken or damaged, the student should report this to dispensary personnel. Refer to the School of Dentistry [Instrument Rental Program User's Guide](#).

SOLICITATION OF PATIENTS

Students who choose to obtain patients through solicitations/requests or other postings on Instagram, Facebook (or other social networking sites), etc., may not use the UTHHealth and/or the UTHHealth logos and may not use UTHHealth, UTHSC-H, University of Texas or UT School of Dentistry names without first obtaining permission from the Office of Legal Affairs and Office of Public Affairs. In the past, such permission has generally not been granted under circumstances such as these. You may not quote prices for the services in the clinic as part of the solicitation/request for patients. Refer to HOOP Policy 8 Use of University Name, Logo or Seal at <https://www.uth.edu/hoop/policy.htm?id=1447864>)

As a student, you may ask persons interested in dental care to contact you, and you may, in your private email response, identify yourself by name and as a UTHHealth Dental Hygiene student and then inform those persons that the treatment is done at the UTSD clinic, and the UTSD clinic sets the fees. You may only identify UTHHealth or the School of Dentistry in your telephone or private email contact, not in the solicitation/request materials/ads/postings, etc. Any questions regarding these guidelines should be directed to the Associate Dean for Patient Care.

When contacting patients/potential patients, use your google voice account to protect your privacy. Per the Texas State Board of Dental Examiners, all correspondence communications, including text messages, are considered part of the patient's record and may be subjected to subpoena! At the School of Dentistry, **texting** to communicate with a patient is **not acceptable**. If a patient texts:

- Respond to the patient and inform them that you will contact them verbally
- Use Google Voice to verbally respond to the patient's text
- Document all correspondence communications in the patient's record under the "Contact Notes"

Violation of these restrictions will subject a student to disciplinary action.

USE OF SOCIAL MEDIA

Students are to adhere to following social media policy and guidelines which can be found at the following links: <https://www.uth.edu/index/social-media.htm> and HOOP Policy 219 Use of Social Media at <https://www.uth.edu/hoop/policy.htm?id=91780f16-961e-4990-be50-8ac8aface6c7>.

Recording Dental Visits for Social Media Posting

Texas is considered a "one-party consent" state, legally allowing recording if one party (including the patient) consents. However, recording may infringe on other patients' HIPAA-protected rights. Therefore, students and patients are **NOT** allowed to record any interactions in our clinical areas. If they are not compliant, Program Directors/GPDs can further reemphasize this policy to them. Otherwise, contact the Office of Patient Care to discuss this policy. Recording for any educational media is excluded and must be done in a contained setting to comply with HIPAA-protected rights.

CLINICAL PATIENTS FOR LICENSURE EXAMINATIONS

Students attempting to secure a patient for examination for licensure **cannot** receive assistance from faculty members in obtaining or assessing the patient. Selection of an appropriate patient is an important factor in the clinical examination. Patient selection is the candidate's responsibility. Candidates are graded on their ability to accurately determine and effectively interpret patient qualification criteria. This is a graded procedure and an integral part of the examination. Patient qualification is the responsibility of the candidate. Therefore, other dental professionals should not pre-qualify patients for the examination per Texas State Board of Dental Examiners clinical licensure guidelines.

SECTION B

**PROFESSIONAL BACKGROUND
INFORMATION**

OATH OF THE AMERICAN DENTAL HYGIENISTS' ASSOCIATION

In my practice as a dental hygienist, I affirm my personal and professional commitment to improve the oral health of the public, to advance the art and science of dental hygiene and to promote high standards of quality care. I pledge continually to improve my professional knowledge and skills to render a full measure of service to each patient entrusted to my care and to uphold the highest standards of professional competence and personal conduct in the interest of the dental hygiene profession and the public it serves.

STUDENT CHAPTER AMERICAN DENTAL HYGIENISTS' ASSOCIATION (SCADHA)

Objectives

The objectives of SCADHA shall be to support the mission of the American Dental Hygienists' Association (ADHA). The ADHA's mission is to improve the public's total health, by advancing the art and science of dental hygiene by ensuring access to quality oral health care; increasing awareness of the cost-effective benefits of prevention; promoting the highest standards of dental hygiene education, licensure, practice and research; and representing and promoting the interests of dental hygienists.

Goals

The goals of the Organization are to:

Provide entry to professional socialization through participation in the organized activities of the Greater Houston Dental Hygienists' Association (GHDHA), Bay Area Dental Hygienists' Association (BADHA), Texas Dental Hygienists' Association (TDHA), and the American Dental Hygienists' Association (ADHA).

- Keep well-in-formed of current and future legislation affecting the dental hygiene profession.
- Educate the public using preventive and therapeutic practices on an individual and group basis.
- Foster life-long learning through a program of expert speakers, arranged by the Vice-President, subject to approval of the officers of the Organization, and advisor(s).
- Promote the dental hygiene profession to laypersons and the dental community through UTSD Orientation and Open House activities; observance of National Dental Hygiene Month; GHDHA, BADHA, TDHA, and ADHA functions.
- Provide for a viable financial base to support the activities of the Organization. These activities will be funded through an annual, non-refundable student chapter of ADHA Professional Fee to be determined jointly by the Director of the Dental Hygiene Program and faculty advisor(s) of the Organization. Such fees may be dispersed for, but not limited to, ADHA membership, social events, continuing education, GHDHA and BADHA component meetings, and the SCADHA/TDHA Annual Session, awards presentations, and installation of officers, subject to advisor(s) approval.

Officers

The officers of SCADHA shall be a senior SCADHA President, Junior and Senior Class Presidents, Junior and Senior Class/SCADHA Vice Presidents, Junior and Senior Class/SCADHA Secretaries, Junior and Senior SCADHA Treasurers. An ADEA delegate will be elected from the Junior Class.

Qualifications

All dental hygiene students with good academic standing (not on probation) in attendance at UTHealth Houston School of Dentistry may be elected to serve in a student organization office. The SCADHA President shall be a member of the senior class. Each class, junior and senior, shall elect a Class President, SCADHA/Class Vice President, SCADHA/Class Secretary, and SCADHA/Class Treasurer.

Nominations and Elections

All senior SCADHA class officers shall be elected into office at the spring meeting (Early April) of their junior year. All junior Class Officers shall be elected into office early in September. The following criteria shall be used when electing officers:

Guidelines for all Candidates and Elections

1. Interested parties must declare their candidacy two to three (2-3) weeks in the UTSD Office of Student and Academic Affairs prior to the election.
2. Prepare and deliver a speech before the members of the organization.
3. Election shall be by secret ballot in the Office of Student and Academic Affairs. The candidate receiving the majority of votes cast shall be declared elected.
4. Officers must maintain a minimum 2.75 GPA with no failures, disciplinary actions or academic probation to remain in office
5. (An officer will be removed from his/her position due to their ineligibility).

Tenure of Office

All officers shall serve for a one-year term, with the exception of ADEA Representative, which is a two-year term commitment. The newly elected senior officers shall serve from May until the following May. The term for the junior officers shall be September-May.

Vacancies

In the event of a vacancy in one of the offices, the candidate receiving the second highest number of votes shall fill the vacancy. In the event there was only one candidate for the office, a special election shall be held.

Officer Duties

A. SCADHA President (DH Senior Student)

The duties of the SCADHA President shall be to:

1. Preside at all SCADHA meetings.
2. Call special meetings.
3. Form committees, appoint committee members and set deadlines for committee work to be reported and/or completed.
4. Act as a liaison between the Greater Houston Dental Hygienists' Association and the SCADHA organization.
5. Preside over the following standing committees:
 - 5.1 Orientation – Organize Orientation Gathering with Advisor(s) and DH Program Director.
 - 5.2 National Dental Hygiene Month Activities
 - 5.3 SCADHA/TDHA Annual Session- prepare school video, hosting, provide volunteers, etc.
 - 5.4 SCADHA Fundraising
6. Submit an annual report of the activities to the SCADHA Advisor.
7. Submit quarterly articles about SCADHA activities for the GHDHA newsletter.

B. Junior and Senior Class Presidents

The duties of the Junior and Senior Class President shall be to:

1. Preside at all Class meetings.
2. Serve on UTSD Student Council (Senior) and sit on the Dean's Council (Junior and Senior).
3. Preside over the following standing committees:
 - 3.1 Fundraising
 - 3.2 Graduation Functions (Senior)

4. Submit an annual report of the activities to the SCADHA Advisor.

C. SCADHA/Class Vice Presidents

The duties of the Junior and Senior SCADHA/Class Vice Presidents shall be to:

1. Preside at all meetings in the absence of the President (SCADHA Senior)
2. Serve on the UTSD Student Affairs Committee (SCADHA Senior).
3. Preside over the following standing committees:
 - 3.1 Welcome breakfast (Senior)
 - 3.2 Obtain guest speakers for Programs/Lunch & Learns (Senior)
 - 3.3 Holiday Party (Junior)
 - 3.4 Programs (Senior)
4. In the event that the SCADHA President's term of office is terminated, the Senior SCADHA Vice President will preside as President for the duration of the office term.
5. Submit an annual report of the activities to the SCADHA Advisor.

D. SCADHA/Class Secretaries

The duties of the Junior and Senior SCADHA/Class Secretaries shall be to:

1. Keep accurate minutes of each Executive Council meeting.
2. Supervise the bulletin board display.
3. Preside over the following standing committees:
 - 3.1 SCADHA/GHDHA community projects
 - 3.2 Awards Ceremony Slide-Show
 - 3.3 SCADHA mentoring program
4. Keep accurate records of the annual reports of offices and committees. Submit an annual report of this office and compile the annual reports of all the Organization's activities to the SCADHA Advisor.

E. SCADHA/Class Treasurers (Senior Class Treasurer serves as SCADHA Treasurer)

The duties of the SCADHA/Class Treasurer shall be to:

1. Maintain accurate records of the financial status of the senior class and SCADHA.
2. Collect and disburse SCADHA funds
3. Correspond with ADHA regarding membership.
4. Preside over the following standing committees:
 - 4.1 Fundraising
 - 4.2 Market-Place /Annual Session
5. Submit an annual report of the activities to the SCADHA Advisor.

F. ADEA Delegate

The duties of the ADEA delegate shall be to:

1. Attend the annual meeting (Funding provided by the Dean's office).
2. Give an oral presentation of the activities of the annual meeting to the members of the Organization.
3. Submit a written report to the Dental Hygiene Program faculty.

The School of Dentistry offers membership in other student organizations. Through these organizations, you will have exposure to community service, leadership, and scholarship opportunities such as the Hispanic Student Dental Association, the Christian Dental Fellowship, Asian American Student Dental Association, Muslim Dental Association, etc. There may be others announced that include dental hygiene membership.

NOTE: All student organizations are required to register their organizations annually at the institutional level through the school Student Affairs Office. For more information, refer to HOOP Policy 110, Student Organizations at <https://www.uth.edu/hoop/policy.htm?id=1448068>.

THE DENTAL HYGIENE PROGRAM PIN


The pin of The Dental Hygiene Program took its design in part from the seal of The University of Texas School of Dentistry at Houston. The University Seal was designed in 1902 based on the Great Seal of the State of Texas. The Dental Hygiene pin was designed in 1957 and was presented to the first graduating class of dental hygienists.

The shape of the pin, a shield, was taken from the center of the University Seal. On the white shield is placed a blue star, blue being symbolic of sincerity. On the star is inscribed "*Disciplina Praesidium Civitatis*." This is translated from Latin to mean, "Education is the Safeguard of Democracy." In the center of the star is an open book that represents an institution of learning. Above and to the right and left of the book are a wreath and branches of olive and live oak.

The pin is worn only by graduates of The University of Texas School of Dentistry at Houston, Dental Hygiene Program.



Dental Hygiene Pin



The University of Texas School of Dentistry Seal

CODE OF PROFESSIONAL ETHICS

The philosophical, practical science of ethics establishes by reason and intelligent observation principles to direct our human conduct. Professional conduct incorporates the knowledge of these principles into practice. The following principles adopted by the 1974 House of Delegates constitute a guide to the responsibilities of the Dental Hygienist.

Each member of the American Dental Hygienists' Association (ADHA) has the ethical obligation to:

1. Provide oral health care utilizing the highest professional knowledge, judgment, and ability.
2. Serve all patients without discrimination.
3. Hold professional patient relationships in confidence.
4. Utilize every opportunity to increase public understanding of oral health practices.
5. Generate public confidence in members of the dental health professions.
6. Cooperate with all health professions in meeting the health needs of the public.

7. Reorganize and uphold the laws and regulations governing this profession.
8. Participate responsibly in this professional association and uphold its purpose.
9. Maintain professional competence through continuing education.
10. Exchange professional knowledge with other health professions.
11. Represent Dental Hygiene with high standards of personal conduct

AWARDS AND HONORS

Academic Achievement Award

This award is given in recognition of those students maintaining the highest GPA throughout their two years in the Dental Hygiene Program.

American Association of Public Health Dentistry

The American Association of Public Health Dentistry (AAPHD) is sponsoring a national recognition award for senior dental hygiene students who have demonstrated a Special Interest / Achievement in Community Dentistry and Dental Public Health.

Colgate Oral Pharmaceuticals Star Award

The Colgate S.T.A.R. award is offered to graduating dental hygiene students who show excellence and commitment to the hygiene profession by:

- Demonstrating true dedication to the profession.
- Exhibiting extraordinary compassion in patient care.
- Displaying enthusiasm and follow-through for community service.
- Demonstrating outstanding patient education and motivation skills.

E-Portfolio Excellence Award

This annual award is sponsored by the Student Chapter of the American Dental Hygienists' Association to a senior student selected by the Dental Hygiene Faculty whose e-portfolio demonstrates significant information about their dental hygiene educational experience; is well organized, personalized and evidence based; clearly provides evidence that criteria were considered to include a variety of artifacts and documents that provide significant irrefutable evidence demonstrating learning, critical thinking, insight, and serious commitment to growth and learning; and is highly professional in appearance and content.

Greater Houston Dental Hygienists' Society Outstanding Professional Leadership Award

The Greater Houston Dental Hygienists' Association (GHDHA) presents this annual award to a senior dental hygiene student for outstanding leadership and professional growth potential. The recipient of the award is chosen by the Awards Committee of GHDHA from a slate of candidates nominated by the SCADHA Advisors in the Greater Houston area. The Professional Leadership Award recipient will receive a plaque and one-year membership to ADHA. In the spring of each year, three to five qualified students are selected as candidates for the honor and awarded to one student from each school. To receive this award, the student must:

- Maintain at least a 3.0 grade point average during the period of dental hygiene academic education.
- Submit a transcript from the dental hygiene program that the student attends. Write a short essay on your post-graduate goals within your professional association.

- Describe what it means to you to be a part of your professional organization.
- List and explain academic achievements and positions in which you have demonstrated leadership, volunteer work, scholarships awarded, special awards or recognition, and the number of times on the Dean's list, etc.
- List any professional monthly meetings or activities in which you participated with documented attendance for each meeting (such as a log of activities attended).
- Request a letter of recommendation from a faculty member at the student's dental hygiene school.

Applicants will submit their award applications to their SCADHA Advisors. SCADHA Advisors will select the top three to five qualified students as candidates and will mail the award applications and supporting documentation to the GHDHA Awards Committee Chair.

The Hu-Friedy Clinical Achievement / Golden Scaler Award

The Hu-Friedy Clinic Award is presented by the faculty of the Dental Hygiene Program to a graduating student who excels as a clinician. The recipient will be one whose clinical judgment and technical skills are judged superior by the faculty and who has assumed professional responsibility and commitment to patient service. This award is sponsored by The Hu-Friedy Manufacturing Company. To be eligible for this award, the student should:

- Receive an "A" in Clinical Practice I, II, III and IV (should not receive an incomplete in any clinical course or clinical case study).
- Complete clinic requirements prior to the last day of clinic.
- Demonstrate a superior clinical ability in proper instrumentation, polishing and have overall rapport with patients.
- Show genuine concern about the oral health of all her/his patients and motivate them towards good oral health.
- Demonstrate professionalism during all phases of patient contact.
- Organize her/his time and utilize clinic time efficiently.

All students who meet the above criteria are eligible for this award regardless of any previous awards they may have achieved. Overall GPA is not a factor in the selection of this award recipient.

Mentor of the Year Award

The Mentor of the Year Award recognizes a senior dental hygiene student who has unselfishly made a positive contribution of time and counsel towards the growth and development of junior dental hygiene students. The candidates(s) are nominated by junior student(s) by filling out an application and submitting it to the SCADHA advisor(s). The Mentor of the Year Award nominees shall be voted on by junior dental hygiene students at the end of April. The nominee with the majority of the votes becomes the recipient of the award.

Procter and Gamble Preventive Dentistry Award

This award is presented to a senior dental hygiene student who demonstrates a commitment to personalized patient instructions for the maintenance of oral health and prevention of disease. To receive this award the student should:

- Show genuine concern for the oral health of each patient.
- Demonstrate exceptional patient education throughout her/his clinic experience.
- Demonstrate consistently an excellent overall knowledge of preventive oral hygiene aids appropriate to individual patient needs.
- Display professionalism during all phases of patient contact.

All students who meet the above criteria are eligible for this award regardless of any previous awards they may have received. Overall GPA is not a factor in the selection of this award recipient.

Sigma Phi Alpha

Sigma Phi Alpha is the national honor society of the Dental Hygiene profession. Component chapters established by schools of dental hygiene are widely distributed throughout the United States. To be elected to Sigma Phi Alpha is an honor and a privilege. The aim of the society is to stimulate high scholarship, professional accomplishment, and greater service to the field of dental hygiene. The membership shall be limited to 10% of the graduating class. Upper 20% of the class are eligible and shall be selected by faculty based on highest academic ranking and excellence in scholarship and leadership. Any student having been on academic probation may not be considered for this award.

Student Teaching Award

This award is presented to a senior dental hygiene student who demonstrates outstanding student and peer teaching with a high potential for a career in dental hygiene education.

UTSD Dental Hygienists' Alumni Association Award

This award is presented to a senior dental hygiene student by the Dental Hygienists' Alumni Association of The University of Texas School of Dentistry at Houston, Dental Hygiene Program. The award honors an outstanding dental hygiene student in recognition of his/her contributions to the dental hygiene profession during his/her tenure as a student at the School of Dentistry.

UTSD Simulation Award

In the spring of 2014, The Society for Simulation in Healthcare and The University of Texas School of Dentistry at Houston (UTSD) created The Society for Simulation in Healthcare Student Award. The award is presented to a graduating dental hygiene student who has demonstrated the highest interest, participation and performance in simulation during their tenure at UTSD.

UTSD Student Award in Interprofessional Education

In 2014, UTSD instituted a new student award in Interprofessional Education (IPE) to recognize a senior dental hygiene student who has "gone the extra mile" in interacting with other healthcare providers/students. To be eligible for this award, the student must:

- Be enrolled in either the DDS, DH or one of the advanced education programs at UTSD.
- Be in good academic standing.
- Have a demonstrated commitment to interprofessional education as evidenced by participation in at least two of the following:
 - UTHealth IPE offering (e.g., the Deans' Honors Colloquium, IPE elective courses or competitions)
 - UTSD IPE offering (e.g., the Holly Hall Retirement Community Rotation)
 - Events or programs offered by a national organization committed to IPE (e.g., The American Society for Bioethics & Humanities)
 - UTSD or UTHealth IPE committee (e.g., the Center for Interprofessional Education (CIPC) Advisory Council)
- Be recommended for the award by a UTSD faculty member or administrator.

UTHEALTH HOUSTON DENTAL HYGIENE ENDOWMENTS AND SCHOLARSHIPS

The University of Texas at Houston Dental Hygiene Program Class of 2003 Endowment Fund

The Dental Hygiene Class of 2003 created history by being the only class ever at the School of Dentistry to establish an endowment while still in school. The Endowment Fund was established in 2003 to assist a senior dental hygiene student in his/her final year of education. Distributions from the endowment shall be awarded to a senior dental hygiene student who exhibits the following criteria:

- Financial Need
- Maintains an academic GPA of 3.0 or higher
- Compassionate
- Highly focused and motivated
- A team player
- An outstanding dental hygiene citizen among faculty, students, and staff
- A future dental hygiene leader
- An ambassador and advocate to the Dental Hygiene Program

The Greater Houston Dental Hygienists' Association Ushma Ramaiya Memorial Scholarship

The Greater Houston Dental Hygienists' Association Ushma Ramaiya Memorial Scholarship was named in memory of Ushma Ramaiya, a 2001 graduate of the UT School of Dentistry Dental Hygiene Program. This scholarship shall be awarded to senior dental hygiene students in good academic standing. Preference will be given to students who have exhibited the characteristics of compassion, focus and motivation, teamwork, leadership, and advocacy of the profession through outstanding community service.

The Shirah May Hall Memorial Scholarship in Dental Hygiene

The scholarship is awarded to a senior dental hygiene student in good academic standing. To be eligible for the scholarship, the student must be in good academic standing. Preference shall be given to students who have exhibited the characteristics of compassion, focus and motivation, teamwork, leadership, and advocacy of the profession.

Gene Stevenson, DDS, Memorial Award in Evidence-based Dentistry - DH

Distributions from the endowment will support scholarships for dental hygiene students who demonstrate excellence in applying knowledge and scientific evidence in clinical care decisions.

UTSD Faculty Scholarships Dentistry - DH

Distributions of these faculty scholarships will support senior dental hygiene students in good academic standing and with financial needs.

Delta Dental Scholarship for Dental and Dental Hygiene Students

The scholarship is awarded to fourth-year pre-doctoral students and senior dental hygiene students. Preference shall be given to students who have shown the most interest, knowledge, and proficiency in pediatric dentistry following the Regents' Rules and Regulations.

Heartland Dental Hygiene Endowed Scholarship Fund

The Caris Dental Hygiene scholarship is awarded to a senior dental hygiene student based on academic merit. The recipient shall display the attributes of a professional capable of entering a dental group practice specifically including teamwork skills, clinical excellence and leadership.

Cloud Dentistry Dental Hygiene Devotion Award

The Cloud Dentistry Dental Hygiene Devotion is about empowering the dental community by acknowledging individuals who have shown steadfast devotion to their craft. The scholarship is awarded to the graduating dental hygiene student who best exemplifies their dedication to dentistry, both throughout their academic work and their long-term plans for making a difference in the industry and with their patients.

Student scholarships

Numerous scholarships are available for dental hygiene students through the UTHealth Office of Student Financial Services and through professional organizations. The following are helpful links:

UTHealth Student Financial Services

<https://www.uth.edu/sfs/contact.htm>

American Dental Hygienists' Association

<http://www.adha.org/scholarships-and-grants>

Texas Dental Hygienists' Association

<http://texasdha.org/scholarships.html>

American Dental Education Association

<http://www.adea.org/studentaward>

SECTION C

GENERAL CLINIC INFORMATION

GENERAL GUIDELINES FOR ERGONOMIC POSITIONING PRACTICE

The prime objective of ergonomic positioning guidelines is to maintain the concepts of work simplification, and provide the greatest degree of comfort, safety, and health to both the patient and clinician.

Position of the Clinician

For the clinician it is essential to:

1. Center body weight on the stool to obtain maximum stability.
2. Keep back straight and shoulders relaxed.
3. Flex at hips so that trunk and thighs form a 60-90-degree angle.
4. Separate knees and feet to width of hips to maintain proper body support,
5. Raise or lower stool to position knees slightly higher than hip level.
6. Position one or both feet flat on the floor with thigh and calf forming a 60-90-degree (one foot may be placed on the rail of the stool).
7. Field of operation should be at elbow level with elbows relaxed and close to the clinician's sides.
8. Hold head erect with only eyes directed downward.
9. Clinician's face should be no closer than 14-16 inches from patient.
10. Care must be taken to never lean on the patient, patient's chair, or rest instruments on the patient's chest.

Patient Positioning

1. Seat the patient with the chair in an upright position.
2. Recline patient so that the patient is in a supine position with head and feet at approximately the same level when working on the maxillary.
3. A semi supine position is used to treat the mandibular arch. The back is slightly raised with the chin down.

Stool to Chair Relationship

Position the stool to permit access to the patient's mouth from the side-front, the side, or the side-back. The hours of the clock can be used to designate zones of operation. The top of the patient's head appears at 12:00 with the feet toward 6:00. More than one operator stool position is determined by a number of factors including:

1. Clinician's height
2. Clinician's arm length
3. Patient's size
4. Cubicle size and arrangement

Patient/Clinician Position for Sextants: Right-handed Clinician

TABLE 2-1. CLOCK POSITIONS—POSITIONING SUMMARY

Treatment Area	Clock Position	Patient Head Position
Mandibular arch—Anterior surfaces toward	8–9	Chin-down; neutral to turned right or left
Maxillary arch—Anterior surfaces toward	8–9	Chin-up; neutral to turned right or left
Mandibular arch—Anterior surfaces away	11–1	Chin-down; neutral to turned right or left
Maxillary arch—Anterior surfaces away	11–1	Chin-up; neutral to turned right or left
Mandibular arch—Posterior aspects toward	9	Chin-down; neutral
Maxillary arch—Posterior aspects toward	9	Chin-up; neutral to turned slightly away
Mandibular arch—Posterior aspects away	10–11	Chin-down; toward
Maxillary arch—Posterior aspects away	10–11	Chin-up; toward

* Patient position always given in relation to the clinician

* When the mirror is not used, the left index finger is generally used to aid with lip or cheek retraction.


Patient/Clinician Position for Sextants: Left-handed Clinician

TABLE 2-2. CLOCK POSITIONS—POSITIONING SUMMARY

Treatment Area	Clock Position	Patient Head Position
Mandibular arch—Anterior surfaces toward	3–4	Chin-down; neutral to turned right or left
Maxillary arch—Anterior surfaces toward	3–4	Chin-up; neutral to turned right or left
Mandibular arch—Anterior surfaces away	11–1	Chin-down; neutral to turned right or left
Maxillary arch—Anterior surfaces away	11–1	Chin-up; neutral to turned right or left
Mandibular arch—Posterior aspects toward	3	Chin-down; neutral
Maxillary arch—Posterior aspects toward	3	Chin-up; neutral to turned slightly away
Mandibular arch—Posterior aspects away	1–2	Chin-down; toward
Maxillary arch—Posterior aspects away	1–2	Chin-up; toward

* Patient position always given in relation to the clinician.

* When the mirror is not used, the right index finger is generally used to aid with lip.



General Principles

1. Posterior sextants - begin with most posterior tooth, and stop at canine.
2. Anterior sextants - from one canine to the other canine.
3. Individual tooth sequence:
 - a. posterior tooth - start on distal line angle to proximal, continue to facial /lingual surfaces and finish on the mesial.
 - b. anterior tooth - start at midline and continue to the proximal surface.
4. The intra oral fulcrum should be:
 - a. inside the mouth
 - b. on the same arch as the tooth being worked on
 - c. as close as possible to the tooth being worked on
 - d. The extra oral fulcrum should be a stabilizing point on the chin or cheek.
5. The stroke will be:
 - a. a light exploratory stroke
 - b. a small, firm working stroke (pull) when removing deposits.

Wrist and arm movements will be used to direct the instrument blade in a vertical direction. The total instrument grasp/hand should pivot from the fulcrum.

Precautions and Variations

In all cases, health is the most important factor and adjustments may be necessary to provide safety and comfort for both patient and operator. To determine correct positioning, the operator must refer to the patient's medical history for pertinent information.

1. A patient with a history of cardiac or respiratory problems may exhibit difficulty in breathing if the chair is fully reclined. The patient would be placed in a more semi supine position.
2. A history of back injury or muscle spasms may require an adjustment in the patient's position to maintain comfort. The patient would be placed in a more semi supine position.
3. For patients with a history of fainting tendency or low blood pressure, ask them to remain seated for a few minutes before getting out of their chair. The sudden movement may cause orthostatic hypotension which causes the patient to become dizzy.
4. When seating an obese patient, use caution when reclining the chair. Heavy weight can stress the backrest.
5. When seating a pediatric patient, have the child slide up in the chair until his head is at the top of the backrest.
6. Pregnant patients in their last trimester may be uncomfortable in a fully reclined position. The patient would be placed in a more semi supine position.
7. If a patient has a sinus condition with post-nasal drainage, the chair should be slightly elevated for comfort.
8. During ultrasonic scaling (without an assistant aid in suctioning), the water collects rapidly in the back of the patient's mouth. Raise the back of the chair slightly for easy water evacuation and patient comfort.

HEALTH HISTORY GUIDELINES

The Patient Interview

When reviewing the health history with a patient who has indicated systemic medical conditions that could be affected by the treatment a dental hygienist would provide, you must be sure to have a thorough understanding of the patient's medical problem and current status of treatment. The information gained from the questions you ask your patients will help you and your instructor plan the appropriate dental treatment for that patient and determine the need for a medical consultation and/or premedication, and appointment accommodations for special needs patients.

Unacceptable Cases

Consultation with a physician [or obtaining medical clearance](#) may be required in some cases. You should use the health history questionnaire and patient interview to identify cases that are not acceptable in the dental hygiene clinic. This would include patients who indicate a history of the following:

1. Active herpetic lesion (labial, facial, or oral)
2. Contagious skin conditions (impetigo, ringworm, scabies)
3. Head lice
4. Conjunctivitis
5. Elevated oral temperature (in excess of 100 degrees F)
6. Respiratory infections involving inflamed throat and/or elevated temperature
7. Active tuberculosis
8. Viral hepatitis (active cases only)
9. Cardiovascular accidents, cardiac bypass surgery or stroke within the last six months
10. Unstable angina
11. Other contagious conditions or diseases

Medical Consultation

Patients with the following conditions will require a consultation with the assigned Group Practice Director to determine if a medical consultation from the patient's physician is required:

1. Uncontrolled Stage II Hypertension
2. Current anticoagulant therapy
3. Heart surgery other than bypass
4. Other systemic diseases, including cardiac arrhythmias, angina, congestive heart failure, renal and hepatic disease
5. Congenital cardiac defects
6. Surgically constructed systemic-pulmonary shunts
7. Diabetes if the patient **has not had** the condition checked by a physician within the last year
8. Uncontrolled, unstable diabetes mellitus (A1C > 7% or fasting Blood glucose reading greater than 154 mg/dl) and uncontrolled Addison's Disease
9. Tuberculosis, if the condition has been active during the last five years
10. Currently under cancer treatment (including long-term chemotherapeutic drug therapy), i.e. bisphosphonates, interferon treatment
11. Current treatment with anti-cancer chemotherapy including use of chemotherapy drugs for noncancerous conditions, i.e. Methotrexate for rheumatoid arthritis, bisphosphonates for osteoporosis
12. Patients who report history of chemotherapy to determine possible use of bisphosphonates
13. Post-irradiation of the mandible or maxilla with greater than 5,000 rads total dose
14. Renal transplant and hemodialysis
15. Glomerulonephritis or other active renal disorder
16. Patient receiving interferon treatment
17. Patients having had a splenectomy
18. Chronic steroid therapy (over 10 days) within the last two years (20 mg./day)
19. Blood diseases, especially acute leukemia, agranulocytosis, granulocytopenia aplastic anemia and agammaglobulinemia
20. Systemic lupus erythematosus
21. Any immunosuppressed patient such as those with acquired immune deficiency syndrome (AIDS)
22. HIV patients, if patient does not have a reliable disease history
23. High Risk Pregnancy
24. Organ transplant

Resource: **Little and Falace's Dental Management of the Medically Compromised Patient 10th Edition**

Kerr, AR, Miller, C, Rhodus, NL, Stoopler, ET, Treister, NS. Little & Falace's Dental Management of the Medically Compromised Patient, 10th Edition, 2024.

PROPHYLACTIC ANTIBIOTIC THERAPY

Patients who are at risk for the development of infective endocarditis and/or infection of a total joint replacement after dentally-induced bacteremia shall be treated using the current [ADA guidelines](#) unless a significant medical reason exists for deviating from these standards.

People with the highest risk for poor outcomes from Infective Endocarditis (IE) may be prescribed antibiotics (IE prophylaxis) prior to certain dental procedures to reduce their risk of developing it. These include procedures that involve manipulation of gingival tissue or the periapical region of teeth, or perforation of the oral mucosa.

Antibiotic prophylaxis is reasonable before the above-mentioned dental procedures for people with heart valve disease who have any of the following:

- Prosthetic cardiac valves, including transcatheter-implanted prostheses and homografts.
- Prosthetic material used for heart valve repair, such as annuloplasty rings, chords or clips.
- Previous IE.

- Unrepaired cyanotic congenital heart defect (birth defects with oxygen levels lower than normal) or repaired congenital heart defect, with residual shunts or valvular regurgitation at the site adjacent to the site of a prosthetic patch or prosthetic device.
- Cardiac transplant with valve regurgitation due to a structurally abnormal valve.
(AHA, 2023) <https://www.heart.org/en/health-topics/infective-endocarditis>

AP for a dental procedure not suggested

Except for the conditions listed above, antibiotic prophylaxis before dental procedures is not recommended for any other types of congenital heart disease.

DOSAGE FOR PROPHYLACTIC USE OF ANTIBIOTICS

For Dental Procedures: Single Dose 30 to 60 Minutes Before Procedure (American Heart Association, 2021)

Antibiotic Prophylactic Regimens for Dental Procedures			
Situation	Agent	Regimen – Single Dose 30 - 60 minutes before procedure	
		Adults	Children
Oral	Amoxicillin	2 g	50 mg/kg
Unable to take oral medication	Ampicillin OR Cefazolin or ceftriaxone	2 g IM or IV* 1 g IM or IV	50 mg/kg IM or IV 50 mg/ kg IM or IV
	Cephalexin**† OR	2 g	50 mg/kg
Allergic to penicillins or ampicillin – Oral regimen	Doxycycline OR	100 mg	<45 kg, 2.2 mg/kg >45 kg, 100 mg
	Azithromycin or clarithromycin	500 mg	15 mg/kg
Allergic to penicillins or ampicillin and unable to take oral medication	Cefazolin or ceftriaxone†	1 g IM or IV	50 mg/kg IM or IV
*IM – Intramuscular; IV – intravenous ** Or other first or second generation oral cephalosporin in equivalent adult † Cephalosporins should not be used in an individual with a history of anaphylaxis, angioedema or urticaria with penicillins or ampicillin. Clindamycin is no longer recommended for antibiotic prophylaxis for a dental procedure.			

Management of patients with prosthetic joints undergoing dental procedures

Clinical Recommendation:

In general, for patients with prosthetic joint implants, prophylactic antibiotics are **not** recommended prior to dental procedures to prevent prosthetic joint infection.

For patients with a history of complications associated with their joint replacement surgery who are undergoing dental procedures that include gingival manipulation or mucosal incision, prophylactic antibiotics should only be considered after consultation with the patient and orthopedic surgeon.* To assess a patient's medical status, a complete health history is always recommended when making final decisions regarding the need for antibiotic prophylaxis.

Clinical Reasoning for the Recommendation:

- There is evidence that dental procedures are not associated with prosthetic joint implant infections.
- There is evidence that antibiotics provided before oral care do not prevent prosthetic joint implant infections.
- There are potential harms of antibiotics including risk for anaphylaxis, antibiotic resistance, and opportunistic infections like *Clostridium difficile*.
- The benefits of antibiotic prophylaxis may not exceed the harms for most patients.
- The individual patient's circumstances and preferences should be considered when deciding whether to prescribe prophylactic antibiotics prior to dental procedures.

Copyright © 2015 American Dental Association. All rights reserved. This page may be used, copied, and distributed for non-commercial purposes without obtaining prior approval from the ADA. Any other use, copying, or distribution, whether in printed or electronic format, is strictly prohibited without the prior written consent of the ADA.

ADA. Center for Evidence-Based Dentistry™

* In cases where antibiotics are deemed necessary, it is most appropriate that the orthopedic surgeon recommend the appropriate antibiotic regimen and when reasonable write the prescription.
Sollecito T, Abt E, Lockhart P, et al. The use of prophylactic antibiotics prior to dental procedures in patients with prosthetic joints: Evidence-based clinical practice guideline for dental practitioners — a report of the American Dental Association Council on Scientific Affairs. JADA. 2015;146(1):11-16.

GUIDELINES FOR MANAGEMENT OF PATIENTS WITH ELEVATED BLOOD PRESSURE

Located on the UTSD intranet: <https://inside.uth.edu/dentistry/docs/mgmtofthehypertensivepatientpolicy.pdf>

This policy document outlines the parameters that guide decisions relative to the care of patients who present with elevated blood pressure (BP). In November 2017, updated clinical practice guidelines were published by the American College of Cardiology and the American Heart Association that define hypertension (HTN) as a BP reading of 130/80 mm Hg or higher. The revised classification system is as follows.

	Values [mm Hg]	Dental Management
NORMAL	< 120 / < 80	No contraindications to dental treatment
ELEVATED	120-129 / < 80	No contraindications to dental treatment
STAGE 1 Hypertension	130-139 or 80-89	No contraindications to dental treatment
STAGE 2 Hypertension	≥140 or ≥90	No contraindications to dental treatment
Upper-level STAGE 2 Hypertension	160-179 or 100-109	*Defer ELECTIVE treatment and refer to physician promptly for evaluation
HYPERTENSIVE URGENCY*	≥ 180* or ≥ 110*	Defer ALL treatment and refer to physician immediately for evaluation

*Non-compliant w/ therapy; re-institute or intensify anti-HTN Rx therapy; treat anxiety prn

*For borderline values, use professional judgment while taking into consideration patient-specific factors as well as the planned treatment procedures.

Guidelines for Measuring Blood Pressure

1. Focus on accurate measurements. These best practices should be followed when measuring a patient's blood pressure in clinic:
 - a. Use a properly calibrated instrument and the correct cuff size
 - b. Do not measure over clothes
 - c. Have the patient sit quietly for at least 5 minutes prior to, and also during, the BP measurement
 - d. Support the arm and make sure the BP cuff is at heart level
 - e. Measure in both arms, if elevated, and consider the higher reading
2. Any patient with an initial blood pressure measurement that is elevated should be re-measured using a manual sphygmomanometer. The first appearance of sound is used to define Systolic BP (SBP). The disappearance of sound is used to define Diastolic BP (DBP).
3. All BP measurements and vital signs taken must be recorded in the patient's electronic health record (EHR) immediately. If these are not recorded for each appointment, the reason must be clearly documented in the EHR.
4. Utilize appropriate stress management protocols. In patients with hypertension who are anxious or fearful, consider use of intraoperative inhalation sedation with nitrous oxide/oxygen.
5. Additional appointment management protocols:
 - a. Avoid rapid position changes of the dental chair to minimize the risk of orthostatic hypotension.
 - b. For patients with BP measurements greater than 140/90, periodic monitoring of BP during treatment, and at the conclusion of the appointment, is advisable.
 - c. Anesthesia considerations: Levonordefrin should be avoided. For patients with BP measurements \geq 180/110, the use of epinephrine should be limited.
6. Capacity to tolerate care:
 - a. Patients with BP measurements below 160/100 may receive any necessary dental treatment.
 - b. For those presenting with BP \geq 160/100, elective dental treatment may be deferred until the BP is brought under better control as confirmed by receipt of a medical clearance from the patient's primary care physician, internist, or cardiologist.
 - c. If urgent or emergency dental treatment is determined to be required, proceed with as limited and conservative treatment procedures as possible to address the chief complaint and/or relieve acute pain unless the BP is confirmed to be \geq 180/110. At this point, no treatment of any type should be performed without a physician consultation.

NOTE: Superficial surgical procedures, including minor oral and periodontal surgery and non-surgical dental procedures, are classified as low risk. Therefore, it appears that the risk associated with most general, outpatient dental procedures is very low.

Follow-up considerations. Encourage healthy lifestyle changes, Rx compliance, and self-monitoring when discussing a patient's level of BP control. Physician follow-up intervals will vary based on the stage of HTN, type of medication(s), level of BP control, and 10-year cardiovascular disease risk assessment.

SIGNIFICANT HYPERTENSION IN CHILDREN

For Children Aged 1–<13 y	For Children Aged ≥13 y
Normal BP: <90th percentile	Normal BP: <120/<80 mm Hg
Elevated BP: ≥90th percentile to <95th percentile or 120/80 mm Hg to <95th percentile (whichever is lower)	Elevated BP: 120/<80 to 129/<80 mm Hg
Stage 1 HTN: ≥95th percentile to <95th percentile + 12 mmHg, or 130/80 to 139/89 mm Hg (whichever is lower)	Stage 1 HTN: 130/80 to 139/89 mm Hg
Stage 2 HTN: ≥95th percentile + 12 mm Hg, or ≥140/90 mm Hg (whichever is lower)	Stage 2 HTN: ≥140/90 mm Hg

REFERENCES

1. Welton, PK, Carey, RM, et al. Hypertension, November 13, 2017. Guideline for the Prevention, Detection, Evaluation and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines.
2. Little, JW, Miller, C, Rhodus, NL. Little & Falace's Dental Management of the Medically Compromised Patient, 9th Edition, 2018.
3. Joseph T. Flynn, David C. Kaelber, Carissa M. Baker-Smith, Douglas Blowey, Aaron E. Carroll, Stephen R. Daniels, Sarah D. de Ferranti, Janis M. Dionne, Bonita Falkner, Susan K. Flinn, Samuel S. Gidding, Celeste Goodwin, Michael G. Leu, Makia E. Powers, Corinna Rea, Joshua Samuels, Madeline Simasek, Vidhu V. Thaker, Elaine M. Urbina, SUBCOMMITTEE ON SCREENING AND MANAGEMENT OF HIGH BLOOD PRESSURE IN CHILDREN; Clinical Practice Guideline for Screening and Management of High Blood Pressure.

ASA PHYSICAL STATUS CLASSIFICATION SYSTEM

Last approved by the ASA House of Delegates on October 23, 2019

The University of Texas School of Dentistry at Houston Department of Diagnostic Sciences has adopted D.F. McCarthy's Physical Evaluation System to assist in categorizing dental patients from the standpoint of medical risk-factor orientation. It is easily adaptable to the needs of private practice.

"The purpose of this system is to quickly and easily place each patient in an appropriate medical-risk category and to thereby provide dental therapy in comfort and relative safety. During the original physical evaluation, the patient is placed in one of our four physical status classes devised by the American Society of Anesthesiologists. The physical status classification then serves as a helpful guide to the level of dental therapy, suggested management, and treatment modification for the medically compromised patient."

ASA Classification	PS Definition	Therapy Modification	Examples, including, but not limited to:
ASA I	A normal healthy patient.	None (stress reduction as indicated)	Fit, nonobese (BMI under 30), a nonsmoking patient with good exercise tolerance.
ASA II	A patient with mild systemic disease	Possible stress reduction and other modifications as indicated.	Current smoker, social alcohol drinker, pregnancy, obesity (30 < BMI < 40), well-controlled DM/HTN, mild lung disease
ASA III	A patient with severe systemic disease	Possible strict modifications; stress reduction and medical consultation prioritized.	Patient with some functional limitation as a result of disease such as poorly controlled DM or HTN, COPD, morbid obesity (BMI ≥40), active hepatitis, alcohol dependence or abuse, implanted pacemaker, moderate reduction of ejection fraction, ESRD undergoing regularly scheduled dialysis, premature infant PCA < 60 weeks, history (>3 months) of MI, CVA, TIA, or CAD/stents.
ASA IV	A patient with severe systemic disease that is a constant threat to life	Minimal emergency care in office; hospitalize for complicated treatment; medical consultation urged.	Patient with functional limitation from severe, life-threatening disease such as a recent (< 3 months) MI, CVA, TIA, or CAD/stents, ongoing cardiac ischemia or severe valve dysfunction, severe reduction of ejection fraction, sepsis, DIC, ARD or ESRD not undergoing regularly scheduled dialysis.
ASA V	A moribund patient who is not expected to survive without the operation. The patient is not expected to survive beyond the next 24 hours without surgery.	Treatment in the hospital is limited to life support only, e.g. airway and hemorrhage management.	A ruptured abdominal aortic aneurysm, massive trauma, and extensive intracranial hemorrhage with mass effect.
ASA VI	A declared brain-dead patient whose organs are being removed for donor purposes		

Classification Guidelines

Any one of which calls for ASA Class 2 placement:

- Present history of allergic rhinitis (hay fever)
- History of any drugs allergy or hypersensitivity
- Pregnancy
- Current smoker, including e-cigarette, pipe, cigars & cigarettes
- Mild Obesity
- History of Hepatitis B that is currently antigen positive
- Anemia
- Extremes of age
- Chronic bronchitis
- Mild to moderate hypertension
- History of arrested pulmonary tuberculosis without disability
- History of corrected congenital heart disease without disability
- History of chronic glomerulonephritis or pyelonephritis without disability
- History of controlled diabetes mellitus without disability
- History of controlled chronic glaucoma
- History of possible attitudinal problems with health care (as negative experiences with prior practitioners)
- History of behavioral problems with health care (as moderate to extreme anxiety)

Many of the preceding diseases or conditions could become Class 3 or 4, depending upon the history and physical examination. Some doctors drop the patient one class if there are two or more diseases, none of which is disabling, for example: a patient with allergic rhinitis, penicillin allergy, and chronic glomerulonephritis could be placed in ASA Class 3 rather than Class 2. This is a judgment decision and is based upon your perception of physical status related to treatment stresses.

Any one of which calls for ASA Class 3 placement:

- Severe diabetes
- Moderate to severe pulmonary disease
- Angina
- Healed myocardial infarction
- Blood dyscrasias
- Moderate to severe hypertension

PREGNANCY

The American Dental Association (ADA) acknowledges that preventive, diagnostic and restorative dental treatment to promote health and eliminate disease is safe throughout pregnancy and is effective in improving and maintaining the oral health of the mother and her child.

Dental Radiography and Pregnancy

- The ADA recommends the use of dosimeters and work practice controls for pregnant dental staff who work with X-rays.
- Studies of pregnant patients receiving dental care have affirmed the safety of dental treatment.
- The American College of Obstetricians and Gynecologists Committee on Health Care for Underserved Women reaffirmed its committee opinion in 2022 stating: "Prevention, diagnosis, and treatment of oral conditions, including dental X-rays (with shielding of the abdomen and thyroid) are safe during pregnancy".

Medications Considered Safe for use During Pregnancy

- Local anesthesia (with or without epinephrine)
- Penicillin
- Amoxicillin
- Cephalosporins
- Clindamycin
- Metronidazole

BLOOD SUGAR LEVELS

Normal A1C levels can range from 4.5-6 percent. An A1C test result between 5.7 percent and 6.4 percent is considered prediabetes. For those patients with diabetes, a target of 7 percent or less is usually preferred. The chart below indicates the A1C level and corresponding average blood sugar levels in milligrams/deciliter.

Note: Uncontrolled, unstable diabetes mellitus (A1C > 7% or fasting Blood glucose reading greater than 154 mg/dl)

A1C level	Estimated Average blood sugar level
5 percent	97 mg/dL
6 percent	126 mg/dL
7 percent	154 mg/dL
8 percent	183 mg/dL
9 percent	212 mg/dL
10 percent	240 mg/dL
11 percent	269 mg/dL
12 percent	269 mg/dL
13 percent	326 mg/dL
14 percent	355 mg/dL

Important questions to ask diabetic patients and note the answers in DHOTEN:

Did you eat breakfast? Lunch? What time today?
 Did you take your medication? What time today?
 What was your blood sugar reading today? Before you ate or after you ate?
 What was your last A1C reading? When was that reading taken?

MEDICAL CONSULTATION

Physical evaluation by the dental team will determine whether or not a medical consultation is indicated. If doubt exists, a consultation with the physician is recommended. The goal of a physical evaluation by the dental team is to determine the ability of the patient to tolerate a specific procedure or series of treatments – **NOT** to diagnose and treat medical problems. The dental team should propose a tentative treatment plan prior to consultation with the physician. The physician will be asked to evaluate the patient, and will either endorse the proposed treatment, or will make recommendations for treatment modifications.

Student Instructions:

1. The medical consult form is located in the Electronic Health Record (EHR)
2. The student, with the patient in the chair, fills out the form.
 - a. Enter pertinent information on the drop-down questions on the medical consultation form:
 - b. Brief introduction of the patient.
 - c. Medical problem of concern.
 - d. Proposed dental treatment-including anesthesia, pre-medication, and other pertinent information.
 - e. Request that the physician evaluate the patient and render an opinion regarding the patient's ability to tolerate the proposed treatment.
 - f. Ask if there are any recommendations or necessary modifications in regards to the proposed treatment.
3. The patient and faculty will sign the form.
4. The student prints the form and gives it to the patient.
5. The patient hand carries it to the physician's office. The patient cannot bring the form back to school. The Medical Consult **must be faxed to patient services at 713-486-4322**. Patient services will scan the form into the electronic health record once received from the physician.
6. The student makes an entry in the Treatment History in the EHR, which documents the consultation request, and includes the physician's name, address, and telephone number if available.

7. Periodically check the “Attachments” tab in the EHR to see if the consultation report has been returned and scanned into the record.
8. Continue with patient treatment according to physician’s recommendations.

ABCDE SYSTEM OF RECOGNIZING MELANOMA

An efficient way to remember common characteristics of melanoma is to think of them in alphabetical order. ABCDE represents asymmetry, border, color, diameter and evolving. A medical doctor or dermatologist will assess characteristics when assessing and diagnosing melanomas.

- A Asymmetry**, because of its uncontrolled growth pattern.
- B Border irregularity**, often with notching.
- C Color** variegation, which varies from shades of brown to black, white, red, and blue depending on the amount and depth of melanin pigmentation.
- D Diameter** greater than 6 mm (which is the diameter of a pencil eraser).
- E Evolving**, change — in size, shape, color, elevation, or another trait, or any new symptom such as bleeding, itching or crusting — points to danger.

Regular skin assessments at home will assist in preventing any life-threatening situations. Skin cancer is not uncommon and can get prevented if detected early and/or cured.

If a mole or lesion is detected, inform the patient not to panic. However, suggest scheduling an appointment for an evaluation.

Student Instructions:

1. Upon evaluation of the EO/IO exam, if a mole is identified, measure the **elevated** mole. Be sure to measure three dimensions and notate in the form. Example: 2x2x2 mole on forehead.
2. If multiple moles on the face and/or neck, do not measure each mole. Indicate in a general manner. Example: Various, localized, scattered, generalized moles on face ranging from 1x1x1-4x3x4.
3. Indicate in DHOTEN under T, “*advised patient to keep watch and/or recommend medical evaluation for any changes in ABCDE’s of identified mole(s)*”.
4. ***DO NOT reference the mole(s) as a LESION, as this word may have a negative connotation to the patient.

GINGIVAL DESCRIPTION

Descriptive Terminology

The degree of severity and distribution of a change should be noted when examining gingiva. When a deviation from normal affects a single area, it can be designated by the number of the adjacent tooth and the surface of the tissue involved, namely, facial, lingual, mesial, or distal.

- A. Severity: Severity is expressed as slight, moderate, or severe.
- B. Distribution
 1. **Localized**- This means that the gingiva is involved only around a single tooth or a specific group of teeth.
 2. **Generalized**- This means that the gingiva is involved around all or nearly all of the teeth throughout the mouth. A condition may also be generalized throughout a single arch.

3. **Marginal**- A change that involves the free or marginal gingiva. This is specified as either localized or generalized.
4. **Papillary**- A change that involves a papilla but not the rest of the free gingiva around a tooth. A papillary change may be localized or generalized.
5. **Diffuse** - When the attached gingiva is involved as well as the free gingiva, it is referred to as a diffuse change. A diffuse condition is most frequently localized, rarely generalized.
6. **Chronic**- Comes on slowly, long duration, painless unless complicated by acute or sub-acute exacerbations.
7. **Acute**- Painful condition that comes on suddenly and is of short duration.

C. Evaluation

1. **Color**- Described as light, regular or dark pink, (normal), red (erythema), bluish red (magenta), coral-pink or other color variations. May include normal pigmentation considering the patient.
2. **Form**- Contour (both marginal and papillary), knife-edge (normal), rounded, blunted, cratered, flattened, bulbous, clefting, festoon.
3. **Density**- Described as stippled (normal), fibrotic, spongy, smooth (shiny edematous).
4. **Attachment**- Note generalized pocket depth and any localized deep pockets.
5. **Bleeding**- Note any upon probing and describe it as slight, moderate, or severe.

CALCULUS CLASSIFICATION

The classification of the patient is determined by the amount and location of calculus. The difficulty of the case may also be a determining factor, (i.e. tenacity of the calculus and periodontal involvement). These are general descriptions of calculus types and variations may exist within each classification.

The classifications are as follows:

Class 0	No calculus present or, localized grainy texture
Class 1	Light calculus Mostly supragingival to 1-2 mm subgingival Localized to buccal max molars and ling mand anteriors
Class 2	In addition to buccal of max molars and ling mand anterior Additional pieces of light to moderate clickable calculus Mostly located on interproximal surfaces premolars or molars
Class 3	Heavy interproximal calculus that is binding Involving 3 surfaces of most teeth including anteriors
Class 4	Heavy interproximal calculus that is binding Involving 3 or 4 surfaces of most teeth May bridge or form continuous rings

AGE CRITERIA

Child	6-11
Adolescent	12-17
Adult	18-49
Geriatric	50+

NOTE: We only see children 6 and above in the clinic.

SPECIAL NEEDS PATIENT

Patients with special needs are defined as those patients whose medical, physical, psychological, or social situations make it necessary to **modify treatment or provide appointment accommodations** in order to deliver dental treatment for them. These individuals include, but are not limited to, people with mental or physical disabilities, complex medical problems, and/or significant physical limitations. A form should be completed in the EHR designating this modification of treatment.

CLASSIFICATIONS OF PERIODONTAL HEALTH AND DISEASES

Health

Periodontal health can be classified in one of two ways:

- ***Health on an intact periodontium*** is used for the patient that has less than 10% of bleeding on probing (BOP) sites, probe depths (PDs) range from 1-3 mm, no evidence of radiographic bone loss (RBL), no recession or clinical attachment loss (CAL), and the gingiva is characterized to be firm, stippled, and resilient without color change.
- ***Health on a reduced periodontium*** is used for the patient with less than 10% BOP sites, PDs range from 1-3 mm, no new RBL (as evidence of monitoring for 1-2 years), existing CAL from factors other than periodontitis (i.e. recession, orthodontics, endodontic lesion, surgery, trauma, clenching/bruxism, etc.); and all modifying and predisposing factors are controlled.

Stable Periodontitis is used for the patient with a history of periodontitis now exhibiting signs of health, which is characterized by the presence of CAL and RBL, probe depths range from 1-4 mm, and controlled modifying or predisposing factors.

Gingivitis

Gingivitis can be either biofilm-induced or non-biofilm-induced. It is important to classify the type of gingivitis. Biofilm (or plaque)-induced gingivitis can be mediated by local or systemic factors or drug-induced. Non-biofilm-induced gingivitis can be due to: genetics, specific infections, inflammatory/immune conditions, reactive processes, neoplasms, endocrine, nutritional or metabolic diseases, traumatic lesions and or gingival pigmentation.

Gingival inflammation can occur with an intact or reduced periodontium. The following are some examples of how to classify gingivitis:

- ***[Biofilm-induced OR Non-biofilm-induced] Gingivitis on an Intact periodontium*** is used for the patient that exhibits the following characteristics: No CAL, no RBL, BOP $\geq 10\%$ of sites ($< 30\%$ sites is localized; $\geq 30\%$ sites is generalized), PDs: 1-4 mm, with uncontrolled modifying or predisposing factors.
- ***[Biofilm-induced OR Non-biofilm-induced] Gingivitis on a Reduced Periodontium*** is used for the patient with no history of periodontitis AND exhibits the following characteristics: presence of CAL due to other factors (not

from periodontitis), presence of RBL, PDs: 1-3 mm, BOP $\geq 10\%$ sites, and uncontrolled modifying and predisposing factors.

Patients with a history of periodontitis now exhibiting signs of gingivitis can be classified in one of two ways:

- **Gingivitis on a Reduced Periodontium with Stable Periodontitis** is used for patients that exhibit the following characteristics: presence of CAL, presence of RBL, PDs: 1-3 mm, BOP $\geq 10\%$ sites, and controlled modifying and predisposing factors.
- **Periodontal Disease Remission** is used for patients that exhibit the following characteristics: presence of CAL, presence of RBL, PDs: 1-4 mm, BOP $\geq 10\%$ sites, and uncontrolled modifying and predisposing factors.

Periodontitis

Periodontitis is the inflammation of the supporting tissues of the teeth—the periodontium. It is characterized by CAL, RBL, periodontal pockets of 4+ mm, and BOP. It is important to note that patients who use or have a history of using tobacco may not exhibit inflammation and BOP as expected; clinical judgment is needed to classify periodontitis in those instances. The onset may be at any age, but is most commonly detected in adults. The prevalence and severity of the disease increases with age.

Periodontitis is initiated and sustained by bacterial plaque, but host defense mechanisms play an integral role in its pathogenesis. The progressive nature of the disease can only be confirmed by repeated examinations. It is reasonable to assume that the disease will progress further if treatment is not provided.

Classification of periodontitis has three parts: Extent (localized vs generalized), Stage (I, II, III, or IV) and Grade (A, B or C). Extent should be based on the teeth with the greatest severity that define the specific stage (*Tonetti, 2018*). Stage is determined by using CAL as the primary parameter, followed by RBL, tooth loss from periodontitis, and then the presence of probing depths of 4+ mm. Stage can be shifted due to case complexity; for example, for a patient who is classified as a Stage II but has short root trunks and furcation involvement, the diagnosis can be shifted to Stage III. Grade can be determined using direct or indirect evidence of progression. Direct evidence requires clinicians to monitor bone loss over the course of 5 years. In an educational setting where patients are not monitored long-term, indirect evidence can be utilized by estimating progression by calculating percentage of bone loss over age. Once the initial grade is established, grade modifiers (poorly controlled diabetes or heavy cigarette smoking or other systemic factors) should be considered to shift from a moderate to rapid progression rate. The tables below outline a summary for finding Stage and Grade (Papapanou, 2018) <https://onlinelibrary.wiley.com/doi/full/10.1111/jcpe.12946#pane-pcw-references>.

Periodontitis stage		Stage I	Stage II	Stage III	Stage IV
Severity	Interdental CAL at site of greatest loss	1 to 2 mm	3 to 4 mm	≥5 mm	≥5 mm
	Radiographic bone loss	Coronal third (<15%)	Coronal third (15% to 33%)	Extending to mid-third of root and beyond	Extending to mid-third of root and beyond
	Tooth loss	No tooth loss due to periodontitis		Tooth loss due to periodontitis of ≤4 teeth	Tooth loss due to periodontitis of ≥5 teeth
Complexity	Local	Maximum probing depth ≤4 mm Mostly horizontal bone loss	Maximum probing depth ≤5 mm Mostly horizontal bone loss	In addition to stage II complexity: Probing depth ≥6 mm Vertical bone loss ≥3 mm Furcation involvement Class II or III Moderate ridge defect	In addition to stage III complexity: Need for complex rehabilitation due to: Masticatory dysfunction Secondary occlusal trauma (tooth mobility degree ≥2) Severe ridge defect Bite collapse, drifting, flaring Less than 20 remaining teeth (10 opposing pairs)
Extent and distribution	Add to stage as descriptor	For each stage, describe extent as localized (<30% of teeth involved), generalized, or molar/incisor pattern			

The initial stage should be determined using CAL; if not available then RBL should be used. Information on tooth loss that can be attributed primarily to periodontitis – if available – may modify stage definition. This is the case even in the absence of complexity factors. Complexity factors may shift the stage to a higher level, for example furcation II or III would shift to either stage III or IV irrespective of CAL. The distinction between stage III and stage IV is primarily based on complexity factors. For example, a high level of tooth mobility and/or posterior bite collapse would indicate a stage IV diagnosis. For any given case only some, not all, complexity factors may be present, however, in general it only takes one complexity factor to shift the diagnosis to a higher stage. It should be emphasized that these case definitions are guidelines that should be applied using sound clinical judgment to arrive at the most appropriate clinical diagnosis.

For post-treatment patients CAL and RBL are still the primary stage determinants. If a stage-shifting complexity factor(s) is eliminated by treatment, the stage should not regress to a lower stage since the original stage complexity factor should always be considered in maintenance phase management.

CAL = clinical attachment loss; RBL = radiographic bone loss.

Source: Papapanou 2018 <https://onlinelibrary.wiley.com/doi/full/10.1111/jcpe.12946#pane-pcw-references>

TABLE 1B Classification of periodontitis based on grades that reflect biologic features of the disease including evidence of, or risk for, rapid progression, anticipated treatment response, and effects on systemic health

Periodontitis grade			Grade A: Slow rate of progression	Grade B: Moderate rate of progression	Grade C: Rapid rate of progression
Primary criteria	Direct evidence of progression	Longitudinal data (radiographic bone loss or CAL)	Evidence of no loss over 5 years	<2 mm over 5 years	≥2 mm over 5 years
	Indirect evidence of progression	% bone loss/age	<0.25	0.25 to 1.0	> 1.0
		Case phenotype	Heavy biofilm deposits with low levels of destruction	Destruction commensurate with biofilm deposits	Destruction exceeds expectation given biofilm deposits; specific clinical patterns suggestive of periods of rapid progression and/or early onset disease (e.g., molar/incisor pattern; lack of expected response to standard bacterial control therapies)
Grade modifiers	Risk factors	Smoking	Non-smoker	Smoker <10 cigarettes/day	Smoker ≥ 10 cigarettes/day
		Diabetes	Normoglycemic/ no diagnosis of diabetes	HbA1c <7.0% in patients with diabetes	HbA1c ≥7.0% in patients with diabetes

Grade should be used as an indicator of the rate of periodontitis progression. The primary criteria are either direct or indirect evidence of progression. Whenever available, direct evidence is used; in its absence indirect estimation is made using bone loss as a function of age at the most affected tooth or case presentation (radiographic bone loss expressed as percentage of root length divided by the age of the subject, RBL/age). Clinicians should initially assume grade B disease and seek specific evidence to shift towards grade A or C, if available. Once grade is established based on evidence of progression, it can be modified based on the presence of risk factors. CAL = clinical attachment loss; HbA1c = glycated hemoglobin A1c; RBL = radiographic bone loss.

Please remember the following:

- Please refer to the latest American Academy of Periodontology (AAP) guidelines for information on periodontal disease classifications.
- Full mouth charting is the minimum expected for all adult cases seen in our clinics, independent of the periodontal diagnosis. For active periodontitis or periodontal maintenance cases, a full periodontal chart is expected at each recall visit to monitor the progression of disease.
- **The classification guidelines offered in this table are suggestive but not definitive.** Faculty members should offer guidance on a case-by-case basis, depending on individual patient risk factors.

PERIODONTAL CHART

1. **BASELINE** - A baseline reading is completed on each new patient 18 years of age or older. A full periodontal charting is done at least once per year in adults, or according to individual patient needs (i.e. aggressive periodontitis in a 15-year-old patient); patients with periodontitis or on a periodontal maintenance recall should have periodontal charting completed at each recall visit. Using the EHR, probe the entire mouth and record all probing depths in the

boxes corresponding to the tooth probed in the row marked PD (probe depths) ... total 6 sites per tooth (3 on Facial and 3 on Lingual).

- a) Mark all bleeding upon probing sites in the row marked BOP (bleeding on probing). Use Y for yes and N for no.
- b) Measure the free gingival margin (FGM) for each tooth by measuring the distance from the CEJ to the free gingival margin, and put numerical value for each tooth in boxes in the row marked FGM-CEJ. It is important to note that the FGM should be measured at the same sites marked for PD measurements. For example, if a 4 mm pocket is found on the DL of #3, then the clinician should first locate the 4 mm to find the corresponding FGM for that distal site. Total of 6 sites per tooth (3 on facial and 3 on lingual—these should correspond to the sites where PD is noted). NOTE: *EHR will calculate the CAL (Clinical Attachment Level) and automatically place it in the CAL row.*
- c) **Tooth mobility** should be recorded in the row of boxes marked MOB using the following classification system (one number per tooth):

Class 0 (0)	No mobility.
Class I (1)	Horizontal movement less than 1mm.
Class II (2)	Horizontal movement greater than 1mm.
Class III (3)	Severe mobility, moves vertically and is depressible in tooth socket.

- d) **Glickman (1953) classifications for furcations.**

Using a Nabers Probe, furcation involvement is usually classified by the amount of a furcation that has been exposed by periodontal bone destruction and is recorded in the FURC row of boxes.

Class I	Early, beginning involvement. A probe can enter the concavity of the furcation area and the anatomy of the roots on either side can be felt by moving the probe from side to side.
Class II	Moderate involvement. Bone has been destroyed to an extent that permits a probe to enter the furcation area but not to pass through between the roots.
Class III	Severe involvement. A probe can be passed between the roots through the entire furcation but not visible upon clinical examination. It is typically apparent on radiographs.
Class IV	Severe involvement. A probe can be passed between the roots through the entire furcation. The furcation is clearly visible upon clinical examination. It is apparent on radiographs.

- 2. **PERIO RE-EVAL**- A periodontal re-evaluation assesses the outcomes of scaling and root planing on an active periodontitis case and must be completed 4-6 weeks after the last quadrant of SRP is completed. During this appointment, a new periodontal chart must be completed (including pocket depths, FGM, bleeding points, mobility, and furcations) and findings should be compared with the baseline measurements recorded in the periodontal chart noted during the comprehensive exam. A summary of changes between the two periodontal charts should be included (CAL, PDs, mobility, furcations and BOP) as well as the gingival description.
- 3. **RECARE/PERIODONTAL MAINTENANCE** - A new periodontal chart must be completed at each recare appointment (including pocket depths, FGM, CAL, bleeding points, mobility, and furcations). Measurements should be compared with baseline measurements to track periodic changes in periodontal health.

CARIES DIAGNOSTIC CRITERIA

Caries detection is not something that is always definite and easily decided. Carious lesions differ greatly. At one end of the spectrum there is the grossly decayed tooth which is easily determined to be carious. At the other end of the spectrum there is the area in an occlusal pit which may or may not show obvious signs of caries. It is at this end of the spectrum where a judgment must be made based on the International Caries Detection and Assessment System (ICDAS) criteria.

The ICDAS concept is that the use of a standardized system, based on best available evidence for detecting early and later stage caries severity, should lead to the acquisition of better-quality information which could then be used to inform decisions about appropriate diagnosis, prognosis, and clinical management of dental caries at both the individual and public health levels.

As well as ICDAS being a coding classification there are simple, standard examination processes employed as part of the system. An important element of the examination is the cleaning of teeth to aid detection since caries forms where there has been plaque stagnation. In addition, the use of compressed air is necessary to reveal the earliest visual signs of caries.

NOTE: An explorer should NOT be used for caries detection. Pressure exerted by the explorer could facilitate further breakdown of enamel that may be able to be remineralized.

To facilitate the use of ICDAS in different settings there is a range of validated tools to select from. The resource, "[Guidelines for Caries Detection \(uth.edu\)](https://uth.edu)" is located under "Links" in the EHR toolbar.

A few general considerations also should be listed relative to these written criteria:

1. Stain or pigmentation alone should not be regarded as evidence of decay as either can occur on sound teeth. Ask yourself: is the lesion soft and light brownish, or hard and dark pigmented?
2. Each subject should be examined in the same manner. An examiner, for example, should avoid the temptation to examine a subject more carefully that appears highly susceptible to dental caries and a person less thoroughly who is relatively free of apparent decay.
3. ***These areas should be diagnosed as sound when there is apparent evidence of demineralization, but no evidence of softness.***

Any written definitions of diagnostic criteria are bound to be interpreted and applied differently by different examiners. Some variation in observational procedures, however, may add strength to one's total knowledge about an agent or a procedure, particularly when the results of different clinical investigators agree.

OCCLUSION–PATIENT EVALUATION


Normal (Ideal) Occlusion

Molar relationship: mesiobuccal cusp of maxillary first permanent molar occludes with the buccal groove of the mandibular first permanent molar.

Malocclusion

Class I: Neutroclusion

Molar relationship: same as Normal, with malposition of individual teeth or groups of teeth.






Class II: Distocclusion

Molar relationship: buccal groove of the mandibular first permanent molar is distal to the mesiobuccal cusp of the maxillary first permanent molar by at least the width of a premolar.

Canine relationship: The distal surface of the mandibular canine is distal to the mesial surface of the maxillary canine by at least the width of a premolar.

Division 1: mandible is retruded and maxillary incisors are protruded.

Division 2: mandible is retruded and one or more maxillary incisors are retruded.



Class III: Mesioclusion

Molar relationship: buccal groove of the mandibular first permanent molar is mesial to the mesiobuccal cusp of the maxillary first permanent molar by at least the width of a premolar.

Canine relationship: The distal of the mandibular canine is mesial to the mesial of the maxillary canine by at least the width of a premolar.

If the distance is less than the width of a premolar, the relationship can be classified as *a tendency toward* that class.


Note tendencies and classify both right and left sides of the mouth.

Other types of malocclusion should be noted as well:

- Open Bite: Note teeth that lack contact with teeth in the opposing arch.
- Posterior Crossbite: Note mandibular posterior teeth that are either facial or lingual to their normal position.

- Anterior Crossbite: Note maxillary incisors that are lingual to the mandibular incisors.
- Edge-to-Edge: Note anterior teeth that occlude on the incisal edge to incisal edge..
- End-to-end: Note molar and premolar teeth that occlude cusp-to-cusp.
- Mid-line deviations: Note the direction of the deviation
- Excessive Overjet: Measure with a periodontal probe the amount of horizontal distance between the labial surface of the mandibular incisors and the incisal edge of the maxillary teeth.
- Underjet: Measure the distance between the lingual of the mandibular incisors and the labial of the maxillary incisors when they are lingual to the mandibular teeth.
- Overbite: Note abnormal overbite if the incisal edges of the maxillary anteriors extend beyond the middle third of the mandibular anteriors.

OCCLUSION OF PRIMARY TEETH



- Mesial Step: distal surface of the mandibular primary second molar is mesial of that of the maxillary.
- Flush/Terminal Plane: The distal surfaces of the mandibular and maxillary second primary molars are on the same vertical plane.
- Distal Step: distal surface of the mandibular primary second molar is distal of that of the maxillary.

ELECTRONIC HEALTH RECORD (EHR)

These are samples of notes for treatment. Students must modify them to represent the treatment provided for that day.

D: Diagnosis

H: Health and dental history

O: Oral hygiene instructions

T: Treatment provided

E: Evaluation- how did the patient do

N: Next visit

EHR DHOTEN treatment templates

These are DHOTEN templates. The student will need to modify the DHOTEN sample according to procedures completed during the clinic session.

D0141 Limited Dental Hygiene Oral Evaluation

DHDx: Clinical findings are consistent with a preliminary DH diagnosis of _____ pending radiographic interpretation.

H: Medical history reviewed with patient. No contraindications to treatment. Vitals: BP: P: R: T:

O: Initial O'Leary score ____%. Initial BOP ____%. Patient states they brush ____ per day; use ____ for interdental hygiene with frequency of ____; use ____ toothpaste and ____ mouthrinse. Gingival Description: _____. Plaque located on _____. Supra/Subgingival calculus present on _____. Patient needs to improve on _____. OHI performed: Described the etiology of _____. Showed pt. areas of _____. Taught ____ technique of ____ using Show-Tell-Do. Patient is ____ (highly, moderately, not) motivated.

T: Performed extra and intra oral exam. EO/IO findings: (WNL or findings noted). Completed dental and periodontal charts. ORA completed. Pre-diagnostic exam was completed by _____. The Needs Assessment exam was completed by Dr. _____. The treatment and alternative options were discussed with the patient, and informed consent signed on _____. Goals of treatment and expected outcomes were discussed.

E: Patient tolerated procedures well (or enter other reactions),

N: Begin non-surgical periodontal therapy (or Start Scaling)

D1110 Adult Prophylaxis & D4346 Prophylaxis in the Presence of Generalized Moderate/Severe Gingivitis (in-process)

DHDx: (Generalized or Localized) _____

H: Medical history reviewed with patient. Changes: _____ No contraindications to treatment. Vitals: BP: P: R: T:

O: Patient states they brush ____ per day; use ____ for interdental hygiene with frequency of ____; use ____ toothpaste and ____ mouth rinse. Gingival Description: _____. Plaque located on _____. Supra/Subgingival calculus present on _____. Patient needs to improve on _____. OHI performed: Described the etiology of _____. Showed pt. areas of _____. Taught ____ technique of ____ using Show-Tell-Do. Patient is ____ (highly, moderately, not) motivated.

T: Performed extra and intra oral exam. EO/IO findings: (WNL or findings noted). Removal of calculus deposits completed (UL, UR, LL, LR) with _____ instrumentation. Continue removal of calculus deposits (UL, UR, LL, LR) quadrants.

E: Patient tolerated procedures well (or ____). Post-Op instructions: Rx non-aspirin analgesic PRN for discomfort and warm salt water rinses due to ____ (slight or moderate) bleeding.

N: Continue prophylaxis ____ quadrant (started this visit)

D1110 Adult Prophylaxis & D4346 D4346 Prophylaxis in the Presence of Generalized Moderate/Severe Gingivitis (final)

DHDx: (Generalized or Localized) _____

H: Medical history reviewed with patient. Changes: _____ No contraindications to treatment. Vitals: BP: P: R: T:

O: Final O'Leary: _____. Patient states they brush _____ per day; use _____ for interdental hygiene with frequency of _____; use _____ toothpaste and _____ mouth rinse. Gingival Description: _____. Plaque located on _____. Supra/Subgingival calculus present on _____. Patient needs to improve on _____. OHI performed: Described the etiology of _____. Showed pt. areas of _____. Taught _____ technique of _____ using Show-Tell-Do. Patient is _____ (highly, moderately, not) motivated.

Home Care Comparison: (Improvement or No improvement) from initial visit.

Gingival Description Comparison: (Improvement or No improvement) from initial visit.

T: Performed extra and intra oral exam. EO/IO findings: (WNL or findings noted). Removal of hard deposits completed _____ (include quadrant started previous visit) with _____ instrumentation. Deplaqued and flossed. Administered fluoride varnish.

E: Patient tolerated procedures well (or _____). Post-Op instructions: Rx non-aspirin analgesic PRN for discomfort and warm salt water rinses due to _____ (slight or moderate) bleeding. Instructions given to the patient regarding diet and home care following fluoride application. Patient goals were met by the following: _____. Expected outcomes were met or not met by the following _____.

N: _____ M recare. Referral to _____ (private DDS, Red, Green, Blue, Yellow, Orange,) Practice for restorative needs.

D4341 Scaling and Root Planing Four or More Teeth/Quadrant or D4342 Scaling and Root Planing One to Three Teeth/Quadrant

DHDx: (Generalized or Localized) _____

H: Medical history reviewed with patient. Changes: _____ No contraindications to treatment. Vitals: BP: P: R: T:

O: Patient states they brush _____ per day; use _____ for interdental hygiene with frequency of _____, use _____ toothpaste and _____ mouth rinse. Gingival Description: _____. Plaque located on _____. Supra/Subgingival calculus present on _____. Patient needs to improve on _____. OHI performed: Described the etiology of _____. Showed pt. areas of _____. Taught _____ technique of _____ using Show-Tell-Do. Patient is _____ (highly, moderately, not) motivated.

T: Performed extra and intra oral exam, (WNL or findings noted). Local anesthesia administered by _____; _____ ml (_____ cartridge(s) of _____, _____ % 1: _____ epi. Completed SRP _____ with _____ instrumentation.

E: Patient tolerated procedures well (or _____). Post-Op instructions: Rx non-aspirin analgesic PRN for discomfort and warm salt water rinses due to _____ (slight or moderate) bleeding.

N: Continue SRP _____ (or Re-evaluation in 4-6 weeks)

D4132 Periodontal Re-evaluation (4-6 Week Re-evaluation)

DHDx: (Generalized or Localized) _____

H: Medical history reviewed with patient. Changes: _____ No contraindications to treatment. Vitals:

BP: P: R: T:

O: Patient states they brush _____ per day; use _____ for interdental hygiene with frequency of _____; use _____ toothpaste and _____ mouth rinse. Gingival Description: _____. Plaque located on _____. Supra/Subgingival calculus present on _____. Patient needs to improve on _____. OHI performed: Described the etiology of _____. Showed pt. areas of _____. Taught _____ technique of _____ using Show-Tell-Do. Pt. given _____ OHI aids.

Home Care Comparison: (Improvement or No improvement) from initial visit.

Gingival Description Comparison: (Improvement or No improvement) from initial visit.

Pt. given _____. OHI aids. Patient is _____ motivated (highly, moderately, not motivated).

T: Performed extra and intra oral exam, (WNL or findings noted). Re-evaluation perio chart completed. Re-evaluated all quadrants and completed removal of calculus with (hand instruments/blended instrumentation). Deplaqued and flossed. Administered fluoride varnish. Instructions given to the patient regarding diet and home care following fluoride application.

Bleeding Level Comparison: (Improvement or No improvement) in overall bleeding points. Initial BOP: _____%; Final BOP: _____%.

Probe Depth Comparisons: Generalized (Improvement or No improvement) in overall PPD. General PPD of _____ mm. Localized PPD of _____ mm in area of # _____. Residual deep PPD _____mm of still present on #_____.

Attachment Level Comparison: Attachment level (with or without) significant changes. Interproximal attachment levels range from _____ to _____ mm.

Recession Comparisons: Recession levels (with or without) significant changes. Recession is in range from _____ to _____mm.

Mobility Changes: (improvement or no improvement) in mobility. Teeth# _____ with class _____ mobility

Plan for follow up of the patient: Grad Perio referral (needed or not needed) for # _____ (deep residual pocket). Graduate periodontics referral form was completed; periodontal faculty signed the form. Situation was explained to the patient. Patient accepts and is willing to proceed with the Grad Perio Referral or Perio maintenance schedule.

E: Patient tolerated procedures well (or _____). Post-Op instructions: Rx non-aspirin analgesic PRN for discomfort and warm salt water rinses due to _____ (slight or moderate) bleeding. Instructions given to the patient regarding diet and home care following fluoride application. Patient goals were met by the following: _____. Expected outcomes were met or not met by the following _____.

N: _____ month Periodontal Maintenance Recare (or Perio Referral). Referral to _____ (private DDS, Red, Green, Blue, Yellow, Orange,) practice for restorative needs.

D4910 Periodontal Maintenance

DHDx: (Generalized or Localized) _____

H: Medical history reviewed with patient. Changes: _____ No contraindications to treatment. Vitals: BP: P: R: T:

O: Patient states they brush _____ per day; use _____ for interdental hygiene with frequency of _____, use _____ toothpaste and _____ mouthrinse. Gingival Description: _____. Plaque located on _____. Supra/Subgingival calculus

present on _____. Patient needs to improve on _____. OHI performed: Described the etiology of _____. Showed pt. areas of _____. Taught _____ technique of _____ using Show-Tell-Do. Patient is _____ (highly, moderately, not) motivated.

T: Performed extra and intra oral exam, (WNL or findings noted). The treatment and alternative options were discussed with the patient and (partial) informed consent signed on _____. Goals of treatment and expected outcomes were discussed. Scaled all quadrants with blended instrumentation to completion. Local anesthesia administered by _____; _____ ml (_____ carpules) of _____ % 1: _____ epi. Deplaqued. Administered fluoride varnish (if treatment planned). Post Op instructions given to the patient regarding restrictions in diet and brushing.

E: Patient tolerated procedures well (or _____). Post-Op instructions: Rx non-aspirin analgesic PRN for discomfort and warm salt water rinses due to _____ (slight or moderate) bleeding. Instructions given to the patient regarding diet and home care following fluoride application. Patient goals were met by the following: _____. Expected outcomes were met or not met by the following _____.

N: _____ month Periodontal Maintenance Recare. Referral to _____ (private DDS, Red, Green, Blue, Yellow, Orange,) practice for restorative needs.

SUPPLEMENTAL FLUORIDE

The most common use of a supplemental fluoride product is for caries prevention. If you determine that supplemental fluoride would be beneficial, consult your instructor. Over-The-Counter products are available through the UTSD Bookstore. Prescription products require a dentist's signature and can be prescribed through the EHR. Documentation must be made in the DHOTEN stating the product and usage instructions given to the patient. <https://dentistry.uth.edu/about/docs/cariesriskmgmtrecsandhomecare.pdf>

Below are listed special cases and their optimum usage:

Orthodontic patients - once a day, preferably at night before going to bed.

Radiation therapy patients (head and neck area) - once a day, preferably at night before going to bed to prevent caries due to decreased salivary flow. Custom trays may be used for application.

Rampant caries - once a day to twice a day, preferably at night before going to bed, depending upon the patient.

Preventive - (child >6 or adult) once a day for best results preferably at night before going to bed.

Cementum hypersensitivity - use of fluoride gel or varnish for controlling root hypersensitivity. Some of the causes of tooth sensitivity result from recession or periodontal surgery exposing cementum.

GENERAL APPOINTMENT SEQUENCE

1. Medical & Dental History
Faculty Check-in
2. Extraoral & Intraoral Examination
Faculty Check-in
3. Oral Hygiene Instruction
4. Dental Hygiene Treatment Procedures
Faculty Check-out

Before the clinic huddle/faculty check-in

Student expectations:

- Patients may be brought into the clinic only if a dental hygiene faculty is on the floor.
- Always open digital radiographs, if available, and reduce the window.
- Upon stepping away from the operator, minimize EHR to maintain HIPAA compliance.
- Review previous medical history and treatment notes.
- Have all of all the necessary codes entered.
 - For the initial appointment, the following treatment codes are needed:
 - Limited DH Exam, D0141
 - Case Complete (Exit Interview), D0013
 - Oral Hygiene Instructions, D1330 (times 3)
 - Pre-diagnostic assessment of patient, D0191, for “*Friends and Family*” patients only
 - Assessment, 00010, required only for patients not assigned to a dental student
 - The codes may have been entered and approved as planned by the facilitator when the patient was assigned.
 - If codes are not entered, it is the responsibility of the student to have the codes approved as planned prior to checking in with faculty.
- Respect HIPAA guidelines: do not discuss dental treatment outside of the clinic operator.
- Seat the patient comfortably; offer to hang coat, purse and/or hat.
- Check the patient in on the EHR.
- Verify the patient has completed and signed the HIPAA forms.
- Review with the patient the following:
 - Name, home address, telephone numbers, emergency contact, race and ethnicity.
 - If any information has changed, go to patient services to update the information.

Initial visit in the dental hygiene clinic

FOR FRIENDS AND FAMILY PATIENTS:

- Acquire headphones from the dispensary and show the UTSD Assessment video.
- Attach a note to the Pre-diagnostic assessment of patient code indicating patient watched and understood the video.

FOR ALL PATIENTS:

- *Discuss consent to initial treatment, and have the patient sign the consent prior to initiating medical/dental history and taking vitals.*

Full Medical/Dental History Assessment:

- Open the **Full Medical/Dental History** form in EHR.
- Take and record blood pressure, pulse, and respiration, and temperature in the medical history form as baseline.
- Add a DHOTEN note: record the diagnosis and all vitals in EHR.
- Interview the patient and make comments on all “yes” responses in the **Full Medical/Dental History** form.
- Fill out the *Accommodation* tab, if applicable.
- Complete the *Sleep Questionnaire* tab for all patients.
- Enter all medications in eRx, including all over-the-counter and non-prescription supplements or medications.
 - If the patient does not take any medications, the *Medications* tab should say “No current medications,” and the date of review should be updated to the current date.
- Have the patient sign **Full Medical /Dental History** when complete.

- Present the Medical History to the dental hygiene instructor.
- The dental hygiene instructor will approve all tabs of the **Full Medical/Dental History** form using their ID badge.

Extraoral & Intraoral Examination:

- Offer gauze squares to the patient for the removal of lipstick or lip balm.
- Place a patient napkin across the patient's chest.
- Ask the patient to remove any removable dental prosthesis.
 - Place the prosthetic in a plastic zip bag.
 - Proceed to DH alcove to clean prosthesis in the ultrasonic bath.
- Ask the patient to rinse with mouthwash.
- Lubricate the patient's lips with cocoa butter.
- Open the **DH Extraoral/Intraoral Exam** form and complete all evaluations.
- Open the **DH Clinical Form**, evaluate the prophy classification and complete the *Calculus* tab.
- The dental hygiene instructor will evaluate prophy classification and approve the *Calculus* tab on the **DH Clinical Form**.
- Present **DH Extraoral/Intraoral Exam** to the dental hygiene instructor.
- The dental hygiene instructor will approve the **DH Extraoral/Intraoral Exam** and **DH Clinical Form**, calculus tab, using their ID badge.

For Friends & Family Patients Only – Pre-Diagnostic Exam:

- Complete the **Adult Assessment** form.
- Have the Group Practice Director (GPD) or appropriate dental faculty complete the Pre-diagnostic exam.
- Enter appropriate radiograph codes and write the exam note.
- The dental faculty completing the pre-diagnostic exam will approve the note, the radiograph codes as planned, and approve the **Pre-Diagnostic Assessment** code (D0191).
- Check in with the dental hygiene instructor before proceeding.
- At this point, the student will continue the DH Limited Exam or proceed to take radiographs.

Patient Education:

- *NOTE: In order to improve patient outcomes, OHI will be delivered to every patient, every clinic session.*
- Disclose the patient, record the O'Leary score in the Perio Chart, label the chart as **Plaque/Bleeding Index**. For most patients, only the 1st and final O'Leary score is recorded.
- Perform appropriate OH education based on observations and DH assessment.

Assessment, Diagnosis & Treatment Planning:

- Complete the Dental Chart, indicating existing items and clinically observed items.
- Refer to the radiographs and the **Guidelines for Caries Detection** (available in Axiom).
- Complete the Periodontal Chart, label the chart as appropriate from the drop-down menu.
- Complete the **Oral Risk Assessment** (ORA).
- Formulate and enter the proposed treatment plan into EHR.
- Once radiographs are complete, present the completed dental chart, periodontal chart, ORA and treatment plan to the dental hygiene instructor.
- The dental hygiene instructor will approve the dental chart items, periodontal chart, and ORA, with their ID badge.
- Enter the periodontal diagnosis into **DH Clinical Form**, Periodontal Classification tab.
- The dental hygiene instructor will approve the **DH Clinical Form**, Periodontal Classification tab with their ID badge.
- *NOTE: In order to proceed through the treatment plan, all radiographs must be interpreted (by a dental student) and a dental examination is required.*
- Request the **Needs Assessment Exam** (00010) or initiation of the **Comprehensive Exam** (D00150).

Comprehensive Oral Examination & Needs Assessment Examination:

- Dental Hygiene treatment cannot continue until the following requirements are met:
 - If the patient is assigned to a dental student:
 - Radiographic interpretation is completed by the dental student.
 - The Comprehensive Oral Exam (COE) is initiated by the assigned student.
 - If the patient is not assigned to a dental student:
 - Radiographic interpretation completed by a dental student from group color practice.
 - The Needs Assessment Exam is completed by the Group Practice Director (GPD).
- *NOTE: The Dental Hygiene instructor may now approve the treatment plan.*
- Discuss the treatment plan with the patient. Include:
 - Dental health problems observed.
 - Nature of the proposed dental hygiene treatment.
 - Potential benefits and risks associated with that treatment.
 - Any alternatives to the proposed treatment, the potential risks and benefits of alternative treatments, including not treating the condition.

Temporary Check-Out:

- Straighten tray table. Have a clean mirror and clean gauze on the bracket tray.
- Ask the dental hygiene instructor for a Stop Check.
- The dental hygiene instructor will examine the hard and soft deposits for errors and then allow dismissal of the patient.
- Re-appoint the patient, return personal items to the patient and escort the patient to the waiting area.
- Complete the treatment notes in EHR and convert the codes of all procedures performed in the clinic during the appointment.
- *NOTE: An OHI code must be planned and completed for each appointment.*
- The instructor will approve the DHOTEN and codes as completed or in progress.
- The instructor will assign a daily grade in EHR and grade the patient care in the Google Spreadsheets and Canvas.

By this point, the following EHR forms should be approved by faculty:

- Medical History/Dental History
- DH Extraoral/Intraoral Exam
- Plaque Index - initial appointment
- Dental Chart - clinical findings only
- Oral Risk Assessment
- Perio Chart
- DH Clinical Form – Prophy Classification and Periodontal Diagnosis
- Treatment plan – if COE is initiated, or Needs Assessment is completed
- Daily Clinic grade and codes completed or initiated for the day
- EHR Notes/DHOTEN

Subsequent appointments in the dental hygiene clinic

Before the Clinic Sessions Begins:

- Assure that planned procedures are entered in the student's schedule and a signed consent form is attached
- If additional codes are needed, inform the instructor at the beginning of the clinic session so all consents may be signed by the patient upon arrival.
- Discuss plans for the clinic session with the dental hygiene faculty.

Medical/Dental History Check-In:

- Interview the patient regarding changes to the medical/dental history since last visit.
- Add a DHOTEN note indicating vitals and any changes to medical history. **Do not make changes in the Medical History form.**
- Changes to medications will be made in eRx.
- REVIEW the medications tab at every visit.
- Notify the instructor when you are ready for a Medical/Dental History check-in.
- Have EHR **Full Medical/Dental History** form open.
- Introduce the patient to the instructor and review pertinent findings and medications with the instructor.

EO/IO Examination Check-In:

- Open EHR to **DH Extraoral/Intraoral Exam** form, and radiographs, if available.
- Perform abbreviated oral examination.
- Record changes in EHR Notes/DHOTEN.
- Have a clean mirror and clean gauze on the bracket tray.
- Introduce the patient to the instructor, stand beside the patient, opposite the instructor; adjust operating light.
- Review patient's x-rays, if available, the dental hygiene faculty will verify EO/IO examinations.
- The dental hygiene instructor will instruct the student to complete O'Leary plaque score and patient education.

Patient Education:

- *NOTE: In order to improve patient outcomes, OHI will be delivered to every patient, every clinic session.*
- Disclose the patient, record the O'Leary score in the Perio Chart, label the chart as **Plaque/Bleeding Index**. For most patients, only the 1st and final O'Leary score is recorded.
- Perform appropriate OH education based on observations and DH assessment.

Proceeding with Dental Hygiene Treatment:

- Prior to initiating scaling/polishing procedures, review with faculty all planned procedures.
- Prior to initiating another procedure, faculty will evaluate completion of the procedures.

Subsequent Temporary Check-Out:

- Straighten tray table. Have a clean mirror and clean gauze on the bracket tray. Ask the instructor for a Stop Check.
- The dental hygiene instructor will examine the hard and soft deposits for errors and then allow dismissal of the patient.
- Re-appoint the patient, return personal items to the patient and escort the patient to the waiting area.
- Complete the treatment notes in EHR and convert the codes of all procedures performed in the clinic during the appointment.
- *NOTE: An OHI code must be planned and completed for each appointment.*
- The dental hygiene instructor will approve the DHOTEN and codes as completed or in progress.
- The dental hygiene instructor will assign a daily grade in EHR and grade the patient care in the Google Spreadsheets and Canvas.

For subsequent appointments, the following EHR forms should be approved by faculty.

- Plaque Index- final appointment
- Daily Clinic grade and approval of codes completed or initiated for the day
- EHR Notes/DHOTEN

Final appointment in the dental hygiene clinic

- 72 hours prior to final appointment, send EHR message to PCC asking to have the record audited.
- Complete Medical History, EO/IO and Patient education as outlined in subsequent appointments.
- Verify completion of all scaling or re-evaluation of previously scaled quadrants.
- Verify completion of all Plaque records and periodontal charts, if applicable.
- Complete de-plaquing (after final scale check or re-evaluation of previously scaled quadrants).
- Apply Fluoride varnish, if appropriate.
- Discuss recare/maintenance with the patient, as well as any post op instruction.
- Verify all codes planned by the dh student are approved as complete, including the case complete code.
- Complete Final Checkout as outlined in subsequent appointments.
- Complete **DH Limited Care Form** (if patient is not referred to a dental student for restorative treatment).

At the completion of the final appointment, the following items should be approved by faculty:

- Final Perio Chart
- Final plaque index
- Daily clinic grade
- Approval of codes completed for the day
- EHR notes/DHOTEN
- Completed **DH Limited Care Form** (if patient is not referred to a dental student for restorative treatment).
- Case Complete Code (D0013) - *not the form*
 - The **DH Case Complete** form will be approved by the facilitator at completion of chart audits.
- After chart audit, send an EHR message to your facilitator regarding the case complete.

PROTOCOL FOR TOPICAL ANESTHETIC

Determine the need for topical anesthesia.

- A. Check medical/dental history for contraindications to topical anesthesia, such as allergies (i.e. amides, ester, PABA, oxymetazoline, any components of drug, etc.), pregnancy, malignant hyperthermia, methemoglobinemia, impaired liver allergy to local/topical anesthetics, or information requiring further investigation, such as syncope or adverse reaction to local/topical anesthetic.
- B. Check medical history for contraindications to epinephrine, such as allergy, cardiac problems, high blood pressure, idiosyncratic reaction(s) to epinephrine, or other conditions requiring clearance from the patient's physician.
- C. Ask patients if they are taking any medications, prescription or non-prescription and check for possible drug interaction with medication(s) they are taking. Non-prescription cold remedies containing an antihistamine can elevate the patient's blood pressure.
- D. Go to the dispensary; get topical anesthetic (if you are working on multiple quadrants remember to get enough for each injection), a syringe, a needle, and carpule(s).
- E. Identify injection sites for application. Dry and isolate the area of injection. Apply a small amount of topical anesthetic with a cotton pellet or swab. If using 20% topical benzocaine, the typical onset is 15-30 seconds so the cotton swab can be removed after 30 seconds. **NOTE: DO NOT OVER-APPLY OR LEAVE ON TISSUE FOR EXTENDED PERIOD OF TIME;** Topical anesthetics have higher incidence of toxicity and allergic reactions due to higher drug concentration compared to local anesthetics.

- F. Make complete, accurate, dated entry in EHR DHOTEN stating the type of topical used (% concentration and drug name), amount used, areas of application (under Treatment provided in DHOTEN) and add a statement for any or no adverse reactions to drug (under Evaluation in DHOTEN).

PROTOCOL FOR ORAQIX® USE IN THE DENTAL HYGIENE CLINIC

1. Check medical history for information contradicting the procedure or requiring further investigation and act accordingly (Oraqix® should not be used in patients with history of congenital or idiopathic methemoglobinemia, allergy or sensitivity to lidocaine and prilocaine and severe hepatic disease).
2. Dental Hygiene faculty will sign a Requisition Form for Oraqix® use. Obtain Oraqix® dispenser, blunt-tip applicator and cartridge of Oraqix® from dispensary. (1 cartridge will be sufficient for most full mouth applications)
3. Remove the blunt-tip applicator from the plastic blister tray, break the seal and remove plastic cover from the cartridge-penetrating end of the cannula. Keep hands away from the exposed cannula during mounting and removal to prevent accidental injuries.
4. Attach the blunt-tip applicator to the tip of the Dispenser.
5. Reset the internal ratchet mechanism before loading the first cartridge. This is accomplished by pressing the mechanism-reset button towards the back end of the body.
6. The air bubble present in the Oraqix® cartridge allows the user to determine if the product is in a liquid or gel form. If the bubble is fixed or moves very slowly, cool the cartridge before use to bring the product back to a liquid form. The cartridge may be loaded into the tip or body of the Dispenser
7. Carefully assemble the body and tip of the Dispenser with the cartridge in place holding the Dispenser in front of you with the tip facing right, rotate the tip sleeve section away from you until locked in place.
8. The applicator tip may be bent to improve access to the periodontal pockets, using the cap. If a greater bend than 45° is desired, a double-bend technique is recommended Note: Do not bend the applicator tip more than once in the same location. Breakage may be more likely if bent at the hub.
9. Hold the Dispenser vertically and observe the transparent portion. The air bubble in the cartridge will be visible and can be removed by depressing the paddle. This will provide a more consistent flow of Oraqix®.
10. Dispense Oraqix® by depressing the paddle. The volume of Oraqix® used per tooth is dependent on the periodontal pocket space. Consult the Oraqix® (2.5%/2.5% lidocaine and prilocaine periodontal gel) package insert for specific dose information. Oraqix® should be applied marginally first before dispensing subgingivally into pockets.
11. Oraqix® is a viscous liquid. Dispensing slowly and evenly works best.
12. When the cartridge is nearly empty, the rubber plunger will be visible in the transparent section of the Dispenser.
13. To reload the Dispenser, first depress the reset button. You will hear the ratchet "click" back into the reset position.
14. Holding the Dispenser in front of you with the tip facing right, rotate the tip sleeve section toward you to unlock the Dispenser tip.
15. Remove the empty cartridge.
16. Insert a new Oraqix® cartridge. A new blunt tip applicator may be used if needed.
17. Reposition the cartridge and tip assembly and lock in place as before.

At the End of Use

1. Carefully remove the blunt tip applicator by re-capping first. Use a one-handed technique to prevent accidental exposure to the contaminated cannula. Dispose of in the same manner as a contaminated dental injection needle by placing it in the Sharps container in your cubicle.
2. Make a note of how much Oraqix® was used before disposing of the cartridge. Remove the used cartridge as described above, and place in the Sharps container in your cubicle.

3. If necessary, wash the surface of the Dispenser to remove any debris, blood or saliva that may be present. Return dispenser in 2 parts to dispensary for sterilization
4. Make a complete, accurate, dated entry in EHR DHOTEN stating the amount of Oraqix® used, areas of application (under Treatment provided in DHOTEN) and add a statement for any or no adverse reactions to Oraqix® (under Evaluation in DHOTEN).

PROTOCOL FOR ARESTIN® USE IN THE DENTAL HYGIENE CLINIC

When indicated by the following criteria, Arestin® may be put into a separate treatment plan after being approved by attending DDS faculty. The DDS will make a note in the EHR that Arestin® was ordered, including the site where the Arestin® is to be placed. The DDS will also need to approve the planned procedure in the treatment plan. The code for Arestin® is 4381 (Chemotherapy per tooth) and the description in the EHR is chemotherapy per tooth. There is a \$0.00 fee for DH students to dispense Arestin® since they are enrolled every year in the Arestin® Student Access Program. The DDS must sign a dispensary “Anesthetic” requisition form and list Arestin® next to the “other” category. Applicators may be obtained from the second-floor dispensary with the signed requisition form. Used plastic applicators can be disposed of in the trash and Arestin® syringes must be returned to the dispensary for sterilization.

Indications for use of Arestin®:

Periodontal Re-evaluation

- After completing a full periodontal charting, probing depths should be compared with baseline measurements taken during the initial exam. All areas with remaining periodontal pockets (4+ mm) should be evaluated for residual calculus and local or systemic factors. Areas with residual calculus need further instrumentation. Arestin® can be placed in localized pockets that range between 5-6 mm after debridement is complete either during the re-evaluation or during the periodontal maintenance appointment.
- Arestin® may be placed at the periodontal re-evaluation or periodontal maintenance appointment if the patient exhibits good oral self-care and if there are localized sites per quadrant that have probing depths of 5-6 mm with bleeding with no signs of local contributing factors (residual calculus, active caries, etc.). A referral to graduate Periodontics should be considered if more than 30% of teeth have refractory sites.

Periodontal maintenance

- Isolated areas that present at the maintenance appointment with 5-6 mm and bleeding may be treated with Arestin following a thorough debridement in patients that exhibit good home care.
- Isolated areas at any subsequent periodontal maintenance appointments may also be treated with Arestin® following a thorough debridement, in patients with good home care.

Contraindications

- Hypersensitivity or allergy to tetracyclines.
- Pregnant or nursing patients.
- Autoimmune syndromes
- Pediatric patients.
- For full list, please use following link: <https://www.arestinprofessional.com/siteassets/pdf/arestin-pi.pdf>

Arestin® Procedure

1. Review medical history for any contraindications for the use of a chemotherapeutic agent.
 - a. Arestin® should not be used in a patient who has a known sensitivity to minocycline or tetracyclines.
 - b. Arestin® should also not be used in pregnant or nursing patients.
2. Explain the rationale to the patient.
3. Have requisition signed by DDS for Arestin® in axiUm. One (1) cartridge is needed for each site to be treated.

4. Arestin will be placed after completion of SRP, at perio re-eval, or at periodontal maintenance as advised by your instructor.
5. Insert the Arestin® cartridge into the handle while exerting slight pressure.
Twist until you feel and hear the cartridge “lock” into place.
6. Should you need to manipulate the cartridge tip to reach difficult areas, gently bend the tip, leaving the blue cap on.
7. Place the cartridge tip into the periodontal pocket, parallel to the long axis of the tooth. Be sure not to force the tip into the base of the pocket.
8. Gently press the thumb ring to express Arestin® powder while withdrawing the cartridge tip away from the base of the pocket.
9. Once delivery is complete, retract the ring and remove the Arestin® cartridge and discard.
10. Patients should be given the following post-op instructions:
 - a. Delay brushing the treated area for 12 hours after treatment with Arestin®.
 - b. Abstain from using interproximal cleaning devices around the treated area for 10 days.
 - c. Avoid foods that could traumatize the gingiva.
11. Indicate in DHOTEN notes areas Arestin was placed and how many cartridges were used.

Arestin® Post-Op Instructions

- Avoid chewing hard, crunchy, or sticky foods for 1 week
- Postpone the use of interproximal cleaning devices around the treated sites for 10 days
- Patients should be advised that some sensitivity is expected during the first week after SRP but they should notify the dentist promptly if ***pain, swelling, itching, swelling, rash, papules, reddening, difficulty breathing, or other signs and symptoms of possible hypersensitivity occur.***

PROTOCOL FOR LOCAL ANESTHETIC

- A. Determine the need for local anesthesia.
- B. Check medical/dental history for contraindications to local anesthesia, such as allergies (i.e. amides, ester, PABA, oxymetazoline, any components of drug, etc.) pregnancy, malignant hyperthermia, methemoglobinemia, impaired liver allergy to local/topical anesthetics, or information requiring further investigation, such as syncope or adverse reaction to local/topical anesthetic.
- C. Check medical history for contraindications to epinephrine, such as allergy, cardiac problems, high blood pressure, idiosyncratic reaction(s) to epinephrine, or other conditions requiring clearance from the patient's physician.
- D. Ask the patient if they are taking any medications, prescription or non-prescription and check for possible drug interaction with medication(s) they are taking. Non-prescription cold remedies containing an antihistamine can elevate the patient's blood pressure.

At this point, you should have determined:

1. Are there any contraindications to local anesthesia for this patient?
2. Is epinephrine contraindicated for this patient?

E. Take vital signs:

- a) If blood pressure is >140/90 in an otherwise healthy appearing patient, repeat the blood pressure in 10 minutes. If still elevated, consult with your instructor. It may be necessary to send a medical consult to the patient's physician.
- b) If blood pressure is 140/90 or lower:

- a. Consult with attending D.D.S. and request a syringe, needle and carpule(s). The anesthesia ordered must be signed by a dentist on a requisition form Request “2% Lidocaine with 1: 100,000 epi” or “2% Lidocaine with 1:200,000 epi”. *IF EPINEPHRINE IS CONTRAINDICATED, REQUEST “Carbocaine 3% Plain” or “Citanest 4% Plain.

***Rule of thumb:** request one carpule per quadrant.

You may want to check with the dental faculty member about the type of anesthesia.)

- b. Go to the dispensary; get topical anesthetic (if you are working on multiple quadrants remember to get enough for each injection), a syringe, a needle, and carpule(s).

***Rule of thumb:**

If you are working in the maxillary arch only, request a short needle.

If you are working in the mandibular arch only, request a long needle.

If you are working in both maxillary and mandibular arches, request a long needle.

- c. Consult with the dental faculty member assigned to the dental hygiene clinic. Be ready for review of the chart with the dental faculty before ordering the anesthesia.

- F. Explain procedure to patient and provide appropriate individualized patient education. (Encourage the patient not to swallow, as this may numb the throat area).

- G. If requested by D.D.S assemble syringe and select site for application of topical anesthetic.

To assembly syringe:

- a. Retract the piston fully
- b. Insert the cartridge – rubber stopper first
- c. Engage the harpoon on aspirating syringes (check before proceeding)
- d. Remove the protective plastic cap from the needle and attach the needle to the syringe
- e. Position the bevel
- f. Remove the colored plastic protective cap and expel a few drops to test for proper flow

- H. Dry and isolate the area of injection. Apply a limited amount of topical anesthetic with a cotton pellet or swab. **Note:** Topical anesthetics have higher incidence of allergic reactions and toxicity due to higher drug concentration.

- I. After anesthesia is provided by D.S or D.D.S begin your work. **Do not discard the needle and carpules** until the end of the appointment. The syringe should be capped using one-hand scoop technique when not in use. If more anesthesia is needed:

- a. If you have at least one full carpule of local anesthetic remaining, find the dental faculty member and tell him/her that your patient needs more anesthesia.
- b. If you do not have enough local anesthesia remaining from the first injection, repeat the above procedure and then find the dental faculty member.
- c. Make complete, accurate, dated entry in EHR DHOTEN as to type and amount of anesthetic delivered and by whom. Add a statement for any or no adverse reactions to anesthetic.
- d. The Dental student or DDS giving the anesthetic must make his or her own chart note for the anesthetic given.

- J. Disassemble the local anesthesia syringe by first removing the carpule by pulling back on the thumb ring and disengaging the carpule from the needle. Next, carefully loosen the needle from the syringe by hand and then use cotton forceps to finish twisting the needle off from the syringe before discarding. Discard needle and carpules in a sharps' container.

CLINIC ASSISTANT ROTATION

STUDENT _____

Faculty Signature _____

DATE _____ AM PM

Faculty (opening clinic), please circle one:

Satisfactory

Unsatisfactory

The following are duties to be performed by the clinic assistants. The supervising faculty member that opened the clinic session will follow up on these procedures to ensure the clinic assistant has completed these tasks before signing this rotation sheet. Any unsatisfactory rotations will result in 0.5 points deducted from final grade.

Students: Please review the list of duties and place a check or initials in blank when completed. If a duty was not applicable for that session, please write N/A in the blank.

_____ Report to DH clinic at 8:15 am if assigned to AM rotation or 1:00 PM if assigned to PM rotation and remain on duty until dismissed by the instructor that opened the clinic.

_____ **Fill the ultrasonic bath with water on Monday AM and wipe down with a paper towel.**

_____ Clean dentures and partials for patients. Supplies and ultrasonic bath in DH Faculty Alcove (2422).

_____ Assist dental hygiene students with dispensary items, charting etc.

_____ Replenish the DH alcove of oral hygiene products. Additional supplies can be found in the DH storage closet and the 3rd-floor dispensary.

- Microscope slides & pH strips are stored in the Periodontics and Dental Hygiene office on the 6th floor
- Organize the supplies in the DH storage room. Keep each item in its designated place. Mark the expiration dates on the boxes of toothpaste and mouthwash, and separate them by the expiration date.
- Let the clinic coordinator know when running low on supplies.

_____ Replenish paper towels and gloves as needed for DH cubicles.

_____ **Empty water out of ultrasonic bath on Thursday PM and wipe down with a paper towel**

_____ Follow-up on checked out equipment from the equipment clipboards and help disinfect/ disassemble and return to the proper storage area. Faculty must sign off on the clipboard prior to returning borrowed equipment to the storage cabinet.

_____ **Verify that the phase contrast microscope has been turned off and that the cover has been replaced. Discard slides and slip covers into the biohazardous sharps container. If saline is low in the bottle, take it to the 2nd floor dispensary and have them refill it with the isotonic solution.**

_____ At the end of the day check that the main unit switch as well as the operative light on all dental chairs occupied by DH students are powered off.

_____ Have this rotation sheet signed by the instructor that opened the clinic and placed it in the clinic coordinator cubby.

PEER EVALUATION ROTATION

STUDENT _____

DATE _____

Faculty Signature _____

Satisfactory

Unsatisfactory

This rotation is designed to provide feedback to students and during performance of various clinical procedures. The Peer Evaluator must be in the clinic on time and stay until dismissed by the assigned faculty.

Instructions:

Peer review consists of observing a classmate perform a procedure and providing immediate feedback on her/his performance. **You must observe 5 procedures each with different students.** Use Section D in your DH Handbook to check how well the procedure is performed. Please be tactful when giving constructive criticism. Place the completed sheet in the clinic coordinator's box in the DH Alcove.

When evaluating your peers consider the following:

- The procedure was performed with complete accuracy (no errors).
- Suggestions for improvement.

Student Observed _____
Procedure Observed _____
Feedback _____

Student Observed _____
Procedure Observed _____
Feedback _____

Student Observed _____
Procedure Observed _____
Feedback _____

SECTION D

CLINICAL PROCEDURES

COMPREHENSIVE PATIENT CARE

Every clinical course has competencies and requirements to meet the course objectives. Along with each clinical course requirements, Comprehensive Patient Care competencies are established to meet CODA standards. They can be completed in clinics I, II, III, or IV with every patient treated. Students are considered competent in providing treatment to patients of all ages, needs, levels of periodontal disease [PVV1], and calculus classifications by successfully completing all Comprehensive Patient Care competencies by the end of spring semester of the senior year. *Prior to graduation* each student will complete at least one Comprehensive Patient Care for the classifications of special needs, adult, adolescent, geriatric, and child; along with, at least a gingivitis, Stage 1, Stage 2, and Stage 3 periodontal disease level.

- Each Comprehensive Patient Care Competency must be completed with a 75 or above to be deemed competent.
- The Comprehensive Patient Care portion of the course grade will be the average of all patients completed during Clinical Practice III.
- These patients must be completed in the UT School of Dentistry, Dental Hygiene clinic. The patient can be a new, re-care patient, or a periodontal maintenance patient.
- Patients completed at outside rotations, dental bay, OMP, and while providing dental hygiene care to dental students' patients will offer *supplemental experiences* with these diverse patient types.
- The successfully passed patient will be counted as a completed patient. This patient may qualify for a personal recall patient.
- Patients completed at outside rotations, dental bay, OMP, and while providing dental hygiene care to dental students' patients will offer *supplemental experiences* with these diverse patient types.

COMPETENCY EVALUATION

Competency evaluations are specific to each course. The competency rubrics will be located in Canvas specific to the clinic level, and updated based on Clinical changes. The grading scale will be dependent on which Clinic level the evaluations are demonstrated. The competency difficulty level increases throughout the curriculum. The competency forms will be at a beginning, developing, and competent level. A 75% or higher pass rate per competency.

YOU MUST SUCCESSFULLY DEMONSTRATE COMPETENCE IN TREATMENT PROCEDURES DESIGNATED AS "COMPETENCY DEMONSTRATIONS" FOR CLINICAL PRACTICE AND CONTINUE TO PERFORM THE PROCEDURES AT A LEVEL OF COMPETENCE IN ORDER TO BE ELIGIBLE FOR GRADUATION.

Competencies that require verbal communication on English speaking patients, unless prior instructor approval has been given for the competency to be attempted on a non-English speaking patient.

ASEPTIC TECHNIQUE /INFECTION CONTROL

Please visit the updated school guidelines on disinfecting the unit, barriers, donning and doffing of PPE, and unit clean up. Masks are highly recommended. Guidelines are frequently updated, please refer to the most recent UTSD protocols on the school's Intranet at <https://inside.uth.edu/dentistry/>.

The student is expected to:

Unit Preparation

1. Turn on the PC and log in to the EHR patient record.
2. Remove extraneous items from the floor and field of contamination.
3. Remove jewelry, 20 second hand wash, put on safety glasses and gloves.

Prior to appointment

1. Flush dental water lines for 30 seconds
 - Three hand pieces- hold up purge level for 30 seconds.
 - Hold the button on the Air Water Syringe for 30 seconds.
 - Flush the suction lines with water for 30 seconds - saliva ejector and high-volume evacuator.
2. Remove 2 germicidal wipes from the canister of EPA-approved hospital disinfectant. Carefully wipe the dental unit, hand piece cords and adapter, chair and counter top, PC keyboard to remove blood and debris. Clean from the cleanest area to the dirtiest. **DO NOT TOUCH OR CLEAN MONITOR SCREEN.**
 - saliva ejector and high-volume evacuator (including controls and cord)
 - light handles, switch and arm
 - tray table
 - air/water syringe and cord
 - hand piece adaptors and cords
 - dental chair, headrest control and arm release button
 - operator chair
 - counter tops and drawer handles
 - Computer, keyboard, mouse, and tray
3. With a new germicidal wipe, repeat the previous cleaning. Wipe the following from the cleanest area to the dirtiest, and allow to dry.
4. Wash hands for one – 20 second washings.
5. Apply barriers to the following (option: clean bare hands or wear gloves):
 - headrest
 - light handles and switch
 - tray table
 - air/water syringe
 - high volume evacuator and saliva ejector
 - PC keyboard – only if computer does not have a white keyboard
 - Backrest of operator stool and lever to raise/lower the stool
6. Place a patient napkin on the rear counter area. Place the following on the napkin.
 - Cocoa butter and leave inside the 2x2 gauze square with a cotton applicator for lubricating the patient's lips.
 - Blue sponge square for cleaning instrument tips during the appointment (remove debris as you work to avoid it from drying. Dried debris is hard to remove during sterilization procedures).
 - Disclosing solution swab
 - Cup with mouth rinse (Patient is to rinse 30 seconds prior to treatment.)
7. Retrieve the instrument cassette from the 2nd floor dispensary. Place an unopened cassette (to be opened when the patient is seated) on the cabinet that is covered with plastic (not where the napkin is placed).
8. Assemble other needed items such as patient education supplies; place them in the appropriate areas.
9. Put on appropriate PPE such as gown, safety glasses, and mask before seeing patients.
10. Perform hand hygiene in front of the patient before starting patient care.

Unit Clean-up

1. Perform hand hygiene.
2. Place a paper towel on a horizontal surface countertop. Doff face shield if used. Clean with a paper towel with soap and water. Place in a clean area to dry.
3. Get a plastic container from the 12 O'clock cabinet, place it on the dental chair, and don gloves.
4. Place properly recapped used needles or other sharps in the nearest Sharps container.
5. Carefully remove gross debris from instrument tips with blue sponge (should be minimal if removing debris as you scale).

6. Place instruments properly into the instrument cassette, close the cassette, and place the cassette into the plastic container. Put the lid on the container and carry to the dirty side of the dispensary.
7. Remember to bring the small bag containing hazardous waste from the operatory and place it into the Biohazard red receptacle at the dispensary.
8. Still wearing disposable gown, protective eyewear and gloves, carry the plastic container to the dirty side of the dispensary. Do not touch anything until you reach a contaminated area, drop your items, return with the container.
9. Return to your cubicle with the plastic container. Place the container on the clean side of the left or right side of the cabinet, wash and dry your heavy-duty gloves, pull-out a sani-wipe and clean the container by cleaning the outside first and then the inside. Wait 3 minutes, repeat the process of disinfecting the container. After 3 minutes, place the container in the 12:00 cabinet with the lid loosely so it can dry.
10. Using gloves, remove barriers and dispose of non-medical waste in trash can.
11. Remove 2 wipes from the canister of EPA-approved hospital disinfectant. Using one wipe in each hand, thoroughly wipe:
 - light handles, top of light, switch, and arm of the light handle
 - computer keyboard, mouse, shelf, and wires [do NOT wipe touchscreen monitor]
 - chair, headrest, and armrests
 - all horizontal surfaces, including countertops and work surfaces, within 6 feet of patient care
 - all cabinet pulls, knobs, and high-touch areas
 - dental unit and delivery unit tray, including all hoses, holders, arms, control buttons, and switches
 - the operator and assistant chairs and levers and any other items that were within 6 feet of patient treatment (e.g. endodontic microscope, external evacuation system, etc.)
 - if a lead apron was used, wipe both sides carefully; touching only the loops only, place apron back to its correct location
 - the outside of the canister of wipes and the spray bottle
12. Flush water lines for 30 seconds (with air/water syringe tip and handpiece attachments). Using a plastic cup, flush the low and high-volume evacuation suction.
13. With gloves on, wash safety glasses with antimicrobial soap, rinse and dry.
14. Place the chair in an upright position, place foot controls at the back of the chair, and place the operator's chair in a minimally disruptive position.
15. Turn off the dental unit.

DOFFING CHECKLIST

1. Perform hand hygiene with sanitizer.
2. Untie the gown at the neck and the back. Grasp the gown from the neck area in the back, being careful to touch the inside of the gown with bare hands, and being careful not to touch the contaminated side of the gown to one's clothes. Sleeves will be turned inside out, and the gown will be rolled. Place the gown in the proper disposal bins. Perform hand hygiene.
3. Remove hair cover and discard (scrub cap to put in brown paper bag). Perform hand hygiene.
4. Remove the mask by pulling away from the face without touching the front of the mask. Touch only the mask straps while removing the mask.
5. Put loupes in carrier
6. Remove shoe covers, if worn
7. Perform hand hygiene with sanitizer
8. Retrieve belongings from cabinet or storage area

Infection Control

1. Practice standard precautions
2. Follow good principles of personal hygiene on a daily basis.

3. Follow proper hand washing guidelines, 20 seconds with soap and water.
4. Keep fingernails short.
5. Wear approved clinic attire (scrubs and gowns, do not wear shirt/sweater over scrubs in clinic area).
6. Do not wear jewelry except a watch when in the clinic.
7. Keep hair securely pinned up and pulled back away from face. Wear a bonnet or scrub cap when using ultrasonic and prophylaxis jets. No messy buns, and no loose hair.
8. Practice proper disinfecting protocol. Do not wear headbands or head coverings that cannot be washed or wiped down. No tied bands.
9. Cover visible tattoos.
10. Wear a face shield and ear plugs when using ultrasonic and air polisher.
11. Verify instruments are sterile.
12. Use appropriate barrier techniques, i.e. gloves, mask, protective eyewear, gown.
13. Remove gloves when leaving the cubicle.
14. Wear gloves when cleaning and disinfecting the cubicle.
15. Follow environmental surface asepsis, i.e. wipe clean/wipe again.
16. Provide a needle cap holder when a needle and syringe are present.
17. Manage and dispose of hazardous waste properly (red biohazard bag), in a biohazard bin located at the check-in area of the dispensary.
18. **Keep forms and documents on the side counter area and covered with plastic** (this is not considered a "contaminated" area).
19. Professionalism- be mindful of what you say, where you say, and who you say to in the clinic area.

COMPLIANCE WITH ALL STANDARDS IN THE INFECTION CONTROL SECTION OF THE SCHOOL OF DENTISTRY CLINIC MANUAL AND COMPLETION OF ANNUAL COMPLIANCE TRAINING IS MANDATORY.

Please follow the most updated CDC and UT guidelines for clinic and infection control.

AIR POLISHING

The student is expected to:

1. Students must sign-in on the check-out sheet (sheets are in a clipboard in the DH Alcove) to use the Cavi-jet or Airflow polishing unit and inform the instructor. Add the unit number on the sign-in sheet.
2. Thoroughly review medical/dental history for information that contraindicates proceeding or will otherwise influence the procedure. (Do not use it on patients who have a severe respiratory illness including patients with COVID-19). Other contraindications include: patients with acute necrotizing ulcerative gingivitis or patients known or suspected of having coronavirus, Hepatitis B, AIDS, tuberculosis, or an HIV positive diagnosis, renal disease, metabolic disorders, patients on diuretics and known infectious diseases).
3. Explain procedure and rationale to the patient, providing individualized patient education. Follow the proper Aseptic technique including the proper PPE for this procedure.
4. Assemble armamentarium:
 - a. air polisher unit/Air Flow polishing unit
 - b. air polishing nozzle
 - c. plastic drape for patient
 - d. protective eye glasses for patient and operator
 - e. paper towels for patient
 - f. face mask and shield for operator
 - g. mouth mirror
 - h. sodium bicarbonate, aluminum trihydroxide powder used on patients with sodium restricted diets or glycine powder
 - i. saliva ejector

- j. HVE
 - k. pre-procedural rinse (essential oil or chlorhexidine-based antimicrobial)
 - l. lubricant for patient's lips with a non-petroleum product (i.e., cocoa butter)
5. Wrap hand piece, unit, and bar grips.
 6. Connect the **BLUE** water line to the water outlet. Make sure to turn on the water connector knob at the attachment port.
 7. Connect the **YELLOW** air line to the air outlet.
 8. Fill the powder chamber with either sodium bicarbonate or aluminum trihydroxide **BEFORE the unit** is turned on. Glycine Powder should only be used in the Air Flow air polishing unit (If the need for more powder is desired during treatment, turn the unit off to relieve pressure in the powder chamber.)
 9. Plug in the unit and turn it on.
 10. Flush lines for two (2) minutes prior to nozzle attachment. (Push the **PURGE** button on the Cavijet unit to perform this function.)
 11. Make sure HVE and saliva ejectors are suction ready to use.
 12. Have the patient use a pre-procedural rinse before using the air polisher.
 13. Drape the patient with a plastic apron and provide with paper towels and safety glasses.
 14. Put on a proper PPE: gown, mask, eyeglasses/ face shield, and gloves, and protective gown and use appropriate aseptic technique.
 15. Adjust patient position to the proper angle.
 16. Utilize the foot pedal in first position for delivery of water for rinsing the teeth and tongue.
 17. Utilize the foot pedal in the second position for delivery of water and air polishing powder for the prophylaxis of the teeth. (When the foot is removed from the pedal, a continuous bleeding of air flows through the hand piece).
 18. Change powder flow rate by rotating the adjustable pointer to **H** for heavy stain, and **L** for light stain (clockwise or counterclockwise) respectively.
 19. Lubricate the patient's lips with a non-petroleum lubricant.
 20. Check cleaning spray. This spray can be contained by "cupping" rather than retracting the patient's lips.
 21. Place 2 X 2 gauze squares on the patient's lip near the working area.
 22. Direct tip of the handpiece nozzle approximately 3 – 4 mm from the tooth surface being cleaned; use direct vision and external fulcrums where possible, use a mirror for illumination as needed.
 23. Center the spray on the middle one-third of the tooth and use a constant circular motion.
 24. Aim nozzle toward the enamel area but not in the sulcus.
 25. Direct handpiece nozzle at an angulation of 60 degrees on anteriors and 80 degrees for posteriors.
 26. The handpiece nozzle is held at 90 degrees when used on occlusal surfaces.
 27. Polish 1-2 teeth for 1-2 seconds with the spray (second position on foot switch) and then rinse the area with water only (first position on foot switch).
 28. Use your hand to cup the patient's cheek to contain the aerosol spray.
 29. Use quick, constant sweeping motions of the hand piece in areas of soft tissue.
 30. Avoid use of air polisher on amalgam, porcelain, composite or highly polished metal restorations.
 31. Avoid use of air polishing on cementum, dentin, or soft tissue.
 32. Use HVE suction continuously.
 33. Check for patient comfort both verbally and visually.
 34. Evaluate the patient to determine that all enamel surfaces are stain and plaque free.
 35. Assess procedures and outcomes and determine ways to improve performance.
 36. At the conclusion of the procedure use the **Purge** button to flush lines for two (2) minutes.
 37. Clean and disinfect cubicle and air polishing unit. Follow the proper aseptic technique to clean and disinfect the unit. Wrap the foot pedal in a plastic bag.
 38. Make sure all parts of the unit are in the bin before returning.
 39. Return the air polishing unit neatly into the storage container and return to Room 2412 (DH Storage Room). Sign out on the check-out sheet.
 40. Make complete, accurate, dated chart entries in EHR.

CARIES RISK

CARIES RISK AND RECOMMENDED THERAPIES For patients 6 years or older

LOW CARIES RISK	MODERATE CARIES RISK	HIGH CARIES RISK
No Caries in last 3 years <ul style="list-style-type: none"> No Xerostomia Low consumption of between meal snacking Has Good Plaque Control Has Optimal Fluoride exposure 	1-2 Caries in last 3 years Has at least 1 of these risk factors: <ul style="list-style-type: none"> Moderate between meal snacking, including sugar beverages 2-3/day Poor Plaque control, BCR 11-30% Sub optimal Fluoride exposure 1 incipient caries/root exposure 	3 or more Caries in last 3 years Has 1 or more of these risk factors: <ul style="list-style-type: none"> Xerostomia High Cariogenic diet, >3 between meal snacking, including sugar beverages Poor Plaque Control, BCR >30% No Fluoride exposure Has 2+ incipient lesions Low Socioeconomic status
RECOMMENDED THERAPIES	RECOMMENDED THERAPIES	RECOMMENDED THERAPIES
NONE	<ul style="list-style-type: none"> OTC Fluoride products <ul style="list-style-type: none"> ACT GelKam Listerine Total Care FL varnish Sealants Xylitol gum Nutritional counseling 	<ul style="list-style-type: none"> Rx Fluoride (approve by DDS) FI varnish Sealants Xylitol gum MI+ Paste Nutritional Counseling Powered Toothbrush
6 month recall	4-6 month recall	3 month recall

Caries Risk Management – Recommendations and Home Care

Low-Caries Risk		
<ul style="list-style-type: none"> Good oral hygiene practices: Brush 2x daily with over-the-counter-fluoridate toothpaste, and floss Dental visit yearly 		
Moderate-Caries Risk		
<ul style="list-style-type: none"> Prescription (5,000 ppm F) toothpaste 2x daily OR Over-the-counter- fluoride toothpaste + flossing + 0.05% fluoride mouthrinse daily at night. Dental visits every six months 		
High-Caries Risk		
<ul style="list-style-type: none"> Fluoride varnish Rinse with baking soda (1 tsp in 125 ml of water) daily Rinse 1 min daily, 7 days a month with 0.12% chlorhexidine gluconate Brush 2x daily with prescription fluoride (5,000 ppm F) Dental visits four to six months 		

⊘ All risk levels – Reduce between meal snacks of fermentable carbohydrates ⊘

Modified from: Featherstone JDB, et al. Caries Management by Risk Assessment (CAMBRA)*: An Update for Use in Clinical Practice for Patients Aged 6 Through Adult. J Calif Dent Assoc 2019;47(1):25-34.

<https://dentistry.uth.edu/about/docs/cariesriskmgmtrecsandhomecare.pdf>

CLEANING REMOVABLE PROSTHESES

The student is expected to:

1. Assemble the armamentarium. Gloves should be worn when handling prostheses.
2. Explain procedure to the patient and provide appropriate individualized patient education.
3. Have the patient remove their prosthesis
 - a. Give the patient a paper towel.
 - b. Provide a private environment for removal.
 - c. Respect the patient's wishes not to talk or be seen without prosthesis.
4. Examine the patient's mouth and the prosthesis.
 - a. Seek consultation regarding ill-fitting prosthesis, ulcerations, inflammation, and cracked or broken prostheses.
5. Take partials/denture(s) to the DH alcove for cleaning.
 - a. Put the patient's name on the outside of the baggie with a Sharpie pen.

- b. Place the prosthesis in a zip-lock bag and add enough of the cleaning agent to completely cover the prostheses. Close the zip-lock bag.
 - c. Place the baggie into the ultrasonic bath for a length of time that removes the stain and/or calculus.
 - d. Repeat the procedure until all debris is removed.
6. Use a denture brush and water to remove any remaining debris from the prosthesis.
 - a. Brush over a sink lined with paper towels or filled with water to prevent breakage.
 - b. Be careful not to bend clasps.
7. Evaluate prosthesis
 - a. No calculus or stain visible
 - b. Outer surfaces smooth in appearance and to patient's tongue
 - c. Absence of all polishing and cleaning agents
8. Rinse the prosthesis and keep it stored in water in a secure place until the end of the appointment.
9. Give the denture brush to the patient for home use.

DENTAL CHARTING

The student is expected to complete the dental charting procedure and enter findings into the EHR.:

1. Open EHR to patient record
2. Open "Chart Add" folder icon
3. Open MiPACS for radiographs
4. Use "Guidelines for Caries Detection" link and radiographs in the EHR
5. Place patient and operator in correct position
6. Use light, mirror, compressed air and radiographs to aid examination
7. Use a cotton tipped applicator as stylus for recording on the monitor screen
NOTE: DO NOT touch the monitor with your gloved hands.
8. Indicate missing teeth on the odontogram
 - a. Right click on missing tooth # and make selection
9. Identify existing conditions
 - a. Click "Findings"
 - b. Choose "Existing Rest/Pro..."
 - c. Select tooth on odontogram
 - d. Click icon on right with 2 teeth to add to odontogram
10. Identify suspicious carious areas and other conditions
 - a. click "Findings"
 - b. Choose "Conditions",
 - c. Select tooth on odontogram
 - d. Click icon on right with 2 teeth to add to odontogram
11. Make a tooth note, (ex. attrition) as appropriate, if description is not available
12. Record findings on proper teeth

DESENSITIZING

The student is expected to:

1. Check medical/dental history for contraindications to the procedure or items requiring further investigation, and act accordingly.
2. Select the appropriate agent
 - Topical Fluoride Varnish (5% NaFl)
 - Relief ACP (5% KNO₃, ACP, & Fl)

Topical Fluoride Varnish (5% NaFl)	Relief ACP (5% KNO ₃ , ACP, & Fl)
cotton rolls	cotton rolls
gauze square	gauze square
bend-a-brush	bend-a-brush
saliva ejector	saliva ejector

3. Assemble the armamentarium
 - cotton rolls
 - gauze square
 - bend-a-brush (included in package of fluoride varnish)
 - saliva ejector
4. Explain procedure to patient and provide individualized patient education:
 - Patient plaque control is the most important issue to convey to help improve hypersensitivity.
 - Discuss trigger foods.
5. Make notes in the DHOTEN of the sensitive area and what product/procedure was used.

DISCLOSING PROCEDURE

The student is expected to:

1. Check medical/dental history for contraindications to the procedure or items requiring further investigation, and act accordingly.
2. Discuss the purpose and procedure of disclosing.
3. Assemble the armamentarium:
 - Disclosing solution swab
 - Mouth mirror
 - Hand mirror
4. Remove excess saliva with a saliva ejector.
5. Remove the swab from the wrapper
6. Hold a swab with a color ring up.
7. Place fingers near the ring.
8. Snap the color ring gently to the side. The liquid will flow to the bottom end of the cotton tip.
9. Apply disclosing solution to the clinical crowns of teeth and gum line.
10. Gently rinse with water and advise patients to expectorate or use suction.
11. Guide the patient in discovering the deposits. Patient education should be completed at this point prior to the removal of deposits. Explain the terms plaque, biofilm, material alba, and food debris.
12. Utilize the phase-contrast microscope, if indicated.
13. Record the plaque index in the patient's dental record.

ETHICS / PROFESSIONALISM / CORE VALUES

The following are some factors that will be considered under ethics and professionalism. The student is expected to demonstrate ethical and professional conduct and judgment when representing UTHealth both within the school setting (clinic and classroom) as well as outside of the school. Examples of expected professional conduct are given but will not necessarily be limited to these examples. There are degrees of unprofessionalism within some of the examples given that may result in a failure for the clinic daily grade at the discretion of the faculty.

Examples of positive professional conduct include:

1. Maintain patient confidentiality.
2. Place the patient's welfare, safety and comfort before oneself when planning and implementing patient care.
3. Concern for the patient's welfare, safety and comfort is prioritized over a quest for achieving a certain grade.
4. Provide treatment in accordance with the treatment plan **after** checking in with supervising faculty.
5. Discuss and review medical history with faculty at the start of each appointment prior to completing any aspect of patient care.
6. Acknowledge and adhere to all medical history alerts.
7. Be prepared with all necessary supplies, instruments and equipment at the beginning of an appointment/procedure.
8. Abide by UTSD clinic policies and regulations.
9. Being open and accepting of feedback provided by faculty to improve individual skills.
10. Maintain physical, mental and emotional composure/attitude in all situations-especially in the presence of patients.
11. Maintain respect, concern and be cooperative toward all students, staff, faculty, and administration.
12. Demonstrate sound clinical judgment by integrating knowledge and skills into patient care commensurate with level of experience.
13. Maintain honesty with faculty members, patients, staff and colleagues.
14. Demonstrate adequate and appropriate communication (Verbal, non-verbal, attitudinal).
15. Provides treatment only when faculty is present and has approved the student to proceed with treatment.
16. Complete all patient records and forms comprehensively and objectively, in a timely manner.
17. Coordinate patient care to assure patient needs are met.
18. Fulfill professional commitments made to community partners and/or service events.

EXTRA / INTRAORAL EXAMINATION

The student is expected to put on gown, mask, glasses, and wash hands for 20 seconds before starting the procedure:

1. Apply methods and prepare materials
 - a. Assemble gauze squares, mirror, and Listerine rinse.
 - b. Explain the purpose and routine order of the examination to the patient.
 - c. Ask the patient to rinse with mouth rinse for 30 seconds, expectorate, and pour any remaining rinse into the sink.
 - d. Without touching the cassette inside, remove the cassette from the sterilization bag. hand sanitizer, and put on gloves.
2. Advise the patient to remove corrective glasses if wearing them. Explain that you are about to begin and ask the patient to indicate if there is any discomfort at any point during the exam.
3. Complete Extra oral Exam - Note NSF if there are no o significant findings.
 - a. Observe from front, noting symmetry of face and neck.
 - b. Have the patient move head from side to side to detect masses or restricted mobility.
 - c. Inspect color and texture of skin on face.
 - d. Inspect eyes and eyelids (opened and closed).
 - e. Palpate occipital nodes bilaterally. Inspect neck for color and texture of skin.
 - f. Palpate pre-& post-auricular nodes bilaterally. Check behind the ears.
 - g. Palpate parotid gland bilaterally.
 - h. Palpate bi-digitally the sub mental and submandibular nodes along the angle of mandible and under chin.
 - i. Palpate bi-digitally along sternocleidomastoid muscles checking for the cervical lymph nodes. Turn head side to side and down.
 - j. Palpate thyroid gland with index finger and thumb. Ask the patient to swallow while palpating.
 - k. Palpate TMJ bilaterally from front. Ask the patient to open and close, noting any irregularities or discomfort.
 - l. Remove gloves and discard once you are done with the examination.

4. Complete intraoral exam
 - a. Ask the patient to wear corrective glasses or safety glasses.
 - b. Put on fresh gloves
 - c. Observe the lips and labial mucosa. Dry both the lips and labial mucosa and then recheck.
 - d. Palpate the lips bi-digitally.
 - e. Observe and palpate the maxillary and mandibular mucobuccal fold.
 - f. Palpate gingiva bi-digitally.
 - g. Retract the cheeks and observe the buccal mucosa. Dry the area and recheck.
 - h. Manipulate duct opening of parotid gland, noting saliva.
 - i. Palpate each cheek bi-digitally or bi-manually.
 - j. Observe the tongue: Wrap with gauze to move from side to side and inspect.
 - Evaluate the dorsal surface of the tongue using the mouth mirror.
 - Evaluate the lateral surface of the tongue using the mouth mirror
 - k. Palpate the entire tongue bi-digitally.
 - l. Observe the ventral surface of tongue, floor of the mouth and the lingual frenum. Dry and recheck all areas. Manipulate the Wharton's duct to check salivary flow.
 - m. Palpate floor of mouth bi-manually.
 - n. Observe and palpate hard and soft palates.
 - o. Inspect oral pharynx and tonsillar region. Depress tongue with mouth mirror.
 - p. Ask the patient to bite down on posterior teeth and check occlusion.
5. Observe deviations from normal and record accurately on the clinical exam form in EHR.
6. It is important to follow a routine order of inspection.

FLOSSING

The student is expected to:

1. Select appropriate type of floss (wax, unwaxed, tape, etc) and the length of material (12-18 inches).
 2. Wind material around the middle or fore fingers of each hand. OR Wind most of the floss around the finger of the least dominant hand. The finger of the other hand will serve as a take-up reel for the used floss. * (Variations are acceptable)
 3. Secure floss/tape with the index finger and thumb of each hand, with a length of $\frac{3}{4}$ to 1 inch between each hand.
 4. Introduce 1 inch of floss interproximally through the contact point with a see-saw motion.
 5. Wrap the floss around the tooth in a "C" shape.
 6. Slide the floss up and down the tooth surface, while holding the material firmly against the proximal surface.
 7. Carry the floss below the gingival margin.
 8. Perform the procedure on the adjacent tooth in the interproximal space by moving from the sulcus to the contact, avoiding the papillary tissue.
 9. Remove the floss by holding the material against one tooth and using a see-saw motion through the contact.
 - a. On the maxilla, the floss is stretched over the thumbs, which guide the floss. Place one thumb on the lingual, and one thumb on the facial side with approximately 1-inch of floss between the thumbs.
 - b. On the mandible, the floss is stretched over the index fingers which guide the floss. Place one thumb or index finger on the buccal, and one thumb or index finger on the lingual side with approximately 1-inch between the fingers.
 10. Reposition and repeat motions, winding used floss around the take-up finger to permit access to a fresh span.
 11. Follow a definite sequence.
 12. Adapt to a patient's ability and preferences, if not harmful.
- Recommend flossing aids as appropriate such as holders, super – floss, etc.

FLUORIDE APPLICATION

The student is expected to:

1. Determine contraindications to fluoride varnish (see below)
2. Explain the benefits of fluoride
3. Describe the application procedure, and obtain the consent of the patient or parent if minor.
 - a. The need for fluoride is based on the patient's risk factors for caries and this should be explained to the patient as part of this process.
 - b. Make notations in the EHR if the patient refuses fluoride.
4. Assemble all necessary supplies.
5. Dry teeth with air. (Gauze may be used to dry teeth by quadrant or sextant.)
6. Mix the varnish with the applicator brush until it is uniform in color.
7. Apply a very thin coat of varnish with a bend-a-brush or cotton tip applicator.
8. Rinse immediately to set the varnish.
9. Suction or have patient expectorate for 30 seconds.
10. Advise the patient not to eat crunchy foods, drink hot liquids, or brush their teeth for 4-12 hours.
11. Make notations in the EHR for fluoride application and post-op instructions given.

CONTRAINDICATIONS TO FLUORIDE VARNISH

- Allergies to Colophonium or other active ingredients in the product being used.
- Do not apply on bleeding gingiva.

IMPLANT PATIENTS

The student is expected to:

1. Review patient assessment data to determine contraindications to treatment or other factors that will influence the procedure.
2. Assemble the appropriate armamentarium:
 - mirror - metal mouth mirror is fine
 - titanium scaler*
 - super floss
 - dental floss
 - auxiliary aids – whatever is appropriate “nylon coated” proxabrush, end tuft brush
 - gauze
 - Fine(pink) proxy paste– to polish implant(s)***Check out a plastic probe and titanium implant instruments from the dispensary.**
3. Use correct patient and operator positioning.
4. Explain the procedure to the patient and present appropriate patient education and psychological support (i.e., home care instructions, why procedure is being done, post-operative instructions, etc.).
5. Implant health should be assessed by reviewing radiographs for any bone loss past the first thread and mobility. If baseline periodontal charting is available, the probe depths should be compared at each recare visit. If mobility is present, the patient should be referred to a periodontist and treatment should not be provided around the implant. Review the patient assessment data and treatment plan assessment procedures to decide which area to scale, and consider the following variables:
 - ability to complete the area
 - patient comfort and acceptance
 - need for tissue conditioning

- patient needs
 - location
6. Use the appropriate type of instruments according to the nature and location of the deposits. (Titanium instruments are recommended to be used on implants.)
 7. Correctly grasp the instrument, establish fulcrum, position cutting edge and activate instrument according to the basic principles of instrumentation. **
 8. Completely scale each tooth and/or implant(s) so that:
 - all surfaces are calculus free
 - gingiva is not bleeding profusely or lacerated
 9. De-plaque teeth and/or implants with appropriate agents to remove plaque. (Implants may be polished with tin oxide or fine (pink) proxy.)
 10. Use appropriate auxiliary aids for complete plaque removal.
 11. Allow the patient to rinse thoroughly with water.
 12. Observe the patient for signs of discomfort and use pain-control techniques as needed to ensure comfort.
 13. Evaluate the procedure and final product to determine that criteria were met and identify ways in which performance could be improved.
 14. Make complete, accurate, dated chart entries in EHR.
 15. Clean up the treatment area and armamentarium.

Refer to *Periodontal Instrumentation* by Gehrig.

INSTRUMENTATION


EXPLORERS

The student is expected to:

1. Grasp (use modified pen grasp).
 - a. Hold the instrument handle with index finger and thumb pads.
 - b. Stabilize with the pad of the middle finger on the instrument shank.
 - c. Maintain contacts between index, middle, and third fingers.
 - d. Place index finger and thumb pads at the junction of handle and shank.
 - e. Maintain handle between second knuckle and "V" of thumb and forefinger.
 - f. Rotate tip between thumb and forefinger when adapting to keep flush side of tip on tooth surface.
 - g. Use light pressure.
2. Fulcrum- the support, or point of rest, on which a lever turns or pivots.
 - a. Establish a fulcrum on a stable tooth, finger, vestibule on gauze or prescribed extra oral.
 - b. Establish an embrasure area, occlusal, or incisal surface.
 - c. Position close to the work area, if possible.
 - d. Use constant, equal pressure.
 - e. Pivot on fingertip for adaptation.
 - f. Move hand (up-down, side-side) when pivoting.
3. Stroke
 - a. Make no independent finger motion.
 - b. Select the correct working end.
 - c. Insert the smallest portion of the tip.

- d. Insert tip at oblique angle to epithelial attachment.
 - e. Insert with tip contacting tooth.
 - f. Maintain the side of the tip on the tooth.
 - g. Keep the terminal shank parallel with the long axis of the tooth.
 - h. Maintain maximum contact of the working end with the tooth (1-2 mm).
 - i. Move tip obliquely or vertically to epithelial attachment.
 - j. Move explorer in the direction the tip is pointed.
 - k. Use short, overlapping strokes
 - l. Use overlapping strokes.
 - m. Cover area from epithelial attachment to margin of gingiva (circumferentially).
4. Technique
 - a. Use systematic sequence.
 - b. Use the correct explorer tip for each surface.
 - c. Choose the explorer type recommended for specific areas or needs.
 - d. Use a mouth mirror for tissue retraction as needed.
 - e. Use a mouth mirror for indirect vision.
 5. Patient-operator positioning
 - a. Assume operator position required for field of operation.
 - b. Position patient for efficient access to the field of operation.

Grasp and Fulcrum Image (Gehrig, J. Periodontal Instrumentation)



PERIODONTAL PROBE

The student is expected to:

1. To use a modified pen grasp and light pressure
2. Establish a stable fulcrum before beginning the strokes.
3. Stroke
 - a. Make minimal independent finger motion.
 - b. Place the lateral surface of the tip on the tooth and slide gently under the gingival margin.
 - c. Place the working end parallel (or as parallel as possible) to the surface being probed.
 - e. Use a 1 mm walking stroke.

- f. Remain in the sulcus and cover the entire area using consecutive strokes.
 - h. Overlap strokes upon reinsertion.
 - i. Insert to epithelial attachment with light pressure.
4. Technique
- a. Use a systematic sequence.
 - b. Use a mouth mirror for indirect vision.
 - c. Maintain correct patient/operator position.
 - d. Read probe markings correctly and mark all 6 areas.

CURETS

The student is expected to:

1. To use the modified pen grasp
2. Establish a stable fulcrum before using the stroke.
3. Stroke
 - a. Select correct working end
 - b. Insert the toe with the blade closed
 - c. Open the blade to 60°-80° for working strokes
 - d. Move in direction toe faces
 - e. Hold side of toe and cutting edge against tooth during: exploratory stroke, and working stroke
 - f. Use short, overlapping strokes
 - g. Roll instrument between thumb and forefinger on line angle to adapt side of toe to tooth
4. Student actions
 - a. Maintain terminal shank handle as close to parallel with long axis of tooth as possible.
 - b. Use oblique, vertical and/or horizontal strokes.
 - c. Have no independent finger motion.
 - d. Apply adequate pressure to remove calculus.
 - e. Use light pressure for exploratory stroke
5. Technique
 - a. Use systematic sequence for scaling individual teeth or quadrants.
 - b. Adapt instrument appropriately:
 - anterior instruments from midline to proximal surface
 - posterior instruments from distal line angle to proximal surface
 - c. Position patient for efficient access to areas.
 - d. Assume operator position as needed for field of operation.

SCALING SICKLE SCALERS

The student is expected to:

1. Use the modified pen grasp.
2. Establish a stable fulcrum before beginning the strokes.
3. Stroke
 - a. Select the correct working end.
 - b. Insert tip with blade closed.
 - c. Move in the direction tip faces.
 - d. Maintain tip and cutting-edge flush with tooth during: insertion; exploratory stroke; and working stroke.
 - e. Use short, overlapping strokes
 - f. Roll instrument between thumb and forefinger on line angle to adapt tip to tooth.
 - g. Apply moderate to heavy pressure to remove calculus.
 - h. Use light pressure for exploratory strokes.
4. Student actions
 - a. Hold the handle as close to parallel with a long axis of tooth as possible.

- b. Use with oblique or vertical strokes.
 - c. Have no independent finger motion
5. Technique
- a. Use systematic sequence for scaling individual teeth or quadrants.
 - b. Adapt instruments:
 - 1) Adapt anterior instruments from midline to proximal surface.
 - 2) Adapt posterior instruments from line angle to proximal surface.
 - c. Position patient for efficient access to areas.
 - d. Assume operator position as needed for field of operation.

INSTRUMENT SHARPENING

(Sharpening Horse Technique)

Uses of the sharpening horse

- Sharp cutting edge without wire edges
- Increased safety
- Instrument longevity
- Maintain original instrument design
- Stabilized fulcrum

Advantages of Sharpening

- Easier calculus removal
- Improved stroke control
- Reduced number of strokes
- Increased patient comfort and satisfaction
- Reduced clinician fatigue
- Preserve and conserve the original shape of the blade


The student is expected to:

1. Assemble armamentarium to include: 2x2 gauze, sharpening stone, Sharpening Horse tool, water or oil depending on the stone used, test stick, safety glasses or loops with light attached.
2. Evaluate sharpness continually while using instruments in the clinical situation; if necessary, obtain another sharp instrument or sharpen the instruments during the procedure.
3. Explain the procedure and provide pertinent, individual education to the patient when sharpening instruments in the operatory.
4. Sharpen instrument utilizing basic sharpening procedures (refer to Canvas Faculty Calibration Course "Sharpening Horse Video") and remove any debris from the instrument.
 - establish the correct angle between stone and cutting edge according to the method being used.
 - maintain the correct angle between stone and cutting edge.
 - utilize proper grasp and stroke.
 - work on a stable work surface with maximum illumination.
 - evaluate the instrument before sharpening to determine if proper contour is present.
5. Test for sharpness before determining if the procedure is complete.
6. Evaluate instrument before the end of sharpening procedure for changes in contour or design features.
7. Use procedures to ensure patient safety and comfort and maximize operator efficiency and effectiveness.
8. Evaluate the procedure and final product to determine ways to improve performance.

PROTOCOL FOR SOPROLIFE INTRAORAL CAMERA

1. A Soprolife intraoral camera may be obtained from the second-floor dispensary with a signed requisition form by DH instructor. Make sure all information is filled out including the patient number.
2. The Soprolife kit will contain a hand-piece, intra-oral tip, and dental barrier.
3. Remove Soprolife intraoral camera from kit. The camera is assembled by the dispensary with the dental barrier and intra-oral tip attached to the hand-piece. Connect the camera cord to the Sopro hand-piece. "Camera" will be displayed on the unit screen when the connection is made.
4. Soprolife operates in three modes. The "daylight mode" will allow the clinician to use the white light for intra-oral images. The "diagnosis mode" aids in detection of caries. The "treatment mode" is used during treatment.
5. Select appropriate mode by using the two buttons on the body of the Soprolife. Button I: pressing on button "I" makes it possible to switch from daylight mode to diagnosis aid mode and inversely. Button II: Pressing on the button "II" makes it possible to switch from daylight mode to treatment aid mode and inversely.
6. Adjust focusing modes. On the hand-piece, there is a rotating ring used to focus in four different modes.

- Extra-oral (Portrait).
- Intra-Oral (1 to 5 teeth).
- I LIFE (cavities and pulp chamber observation)
- Macro (details that cannot be seen with naked eye).

9. If the image is blurry check the dental barrier to see if it is correctly positioned on the camera head.
 - a. Note: The camera head should be placed face down when inserting it into the Sopro dental barrier.
10. Create a photography folder in the X-Ray (MiPACS) EHR. Click the "Capture" icon which will turn green when activated.
11. Slightly touch SoproTouch  or briefly press the footswitch as soon as the desired image appears on the monitor. If an assistant is available, the images may also be captured using the "Capture Image" tab located on the monitor screen. The image is automatically stored, and displayed on the screen.
12. Click "End Session" and delete unwanted images prior to saving captured images. Captured images must be approved by faculty to save images in the EHR.
13. Rename the pictures to match tooth numbers
14. Captured images must be approved by faculty to save images in the EHR.
15. Make notes in the DHOTEN.

Diagnostic Mode

1. The Soprolife diagnosis mode is an aid in the detection of caries. The auto fluorescence technology allows the clinician to detect occlusal or interproximal decay. Soprolife has not been proven to detect incipient caries.
2. When the camera is in diagnosis mode it will display a blue light. The blue light will cause healthy dentin to fluoresce with a green color.
3. Any other color than acid green, light green or blue (according to thickness of enamel) displayed in the image are alert signals.
4. Green/black, bright red, black/red, or gray areas are alert signals. Suspicious areas should be checked for calculus, plaque, and preventative materials that can interfere with caries detection.

At the End of Use

1. Disinfect the Sopro camera with a Caviwipe. Remove the dental barrier from the hand-piece. Re-attach the intra-oral tip prior to returning the camera. Dispose of the blue sterilization wrap.
2. Return the Soprolife camera back to the second-floor dispensary.

MEDICAL AND DENTAL HISTORY

The student will complete a medical/dental history on all patients. A full medical history form is filled on all new patients. Review and update of complete medical history is on all new assessment patients, transferred patients and every 6 months of patient's visit using control R on every question. The type of data collected will dictate proper clinical procedure. The medical history will be reviewed, and updated if necessary, at the beginning of each consecutive appointment.

1. The clinician needs to be in proper PPE) for taking Medical History.
2. Assemble armamentarium prior to seating the patient.
3. Review with the patient the Patient Information, needed for record keeping:
 - a. name, home address, telephone numbers, emergency contact
 - b. If any information has changed, send an EHR message stating changes to your PCC
 - c. Observe the patient to confirm there are no symptoms of cold sore
4. Take blood pressure, pulse, respiration and temperature before starting the competency. Make "D" and "H" notes (of the DHOTEN) in the Treatment History tab of the EHR. Document vital signs in the EHR before reviewing medical history with the instructor.
5. **For the DH1 competency (process):** Review the following categories with the patient (ask questions) in front of the instructor: *Baseline Data, General, HEENT, Cardiovascular, and medication information.*
6. Review/question each category of medical conditions listed. Make notations for affirmative answers in "comments" at the end of each category; double click and use text box and date when the condition was diagnosed, what happened, medications, etc. Add N/A in the text box for those categories where all answers are negative by double clicking.
7. Place medications in the eRx, not in the form. Note in the Treatment History if previously taken medication has been discontinued. Delete medications from the medication tab and add the new medication in the eRx, not into the Medical History. Look up all medications (OTC and herbal included) in Lexi-Comp found in the EHR link and be knowledgeable about pharmacological category, indications for use and contraindications; warnings, precautions and adverse reactions to treatments; drug interactions and dental considerations. Update medication tab at every visit after reviewing the medications.
8. Assess need for medical consultation and /or antibiotic premedication or any medical alert (right click on medical alert and highlight) Note...some medications won't automatically be posted as an alert even if listed on the medical history.
9. Place N/A in spaces that have no answer
10. Determine the ASA classification
11. Review the Dental History page (paying particular attention to any affirmative answers).
12. Make appropriate notations in Treatment History DHOTEN not included under the "H" section.
13. Have the instructor review medical and dental history.

NUTRITIONAL COUNSELING

The student is expected to:

1. Review medical/dental history and Oral Risk Assessment form for information to determine if a caries risk and/or periodontal risk dietary assessment is indicated.
2. Explain to the patient all the procedures and rationale for 24-hour dietary recall. Obtain patient permission to proceed.
3. Add Nutritional Counseling code D1310. Receive faculty approval and obtain patient consent.
4. Complete 24-hour food diary with patient using form in Canvas after Oral Risk Assessment form in EHR has been completed.
5. Analyze the 24-hour food diary and assess the patients' exposures to fermentable carbohydrates. Calculate acid exposure times. Review food diary for any nutritional deficiencies that may put the patient at risk for periodontal disease.
6. Discuss the results with the patient.

7. Counsel the patient using motivational interviewing and open-ended questioning. Assist the patient in setting nutritional goals to decrease their caries and /or periodontal disease risk.
8. Refer the patient to <http://www.choosemyplate.gov> for additional guidance.
9. Make complete and accurate entry in the DHOTEN notes including all nutritional recommendations.
10. Turn D1310 code to "I" in process if not completed at the same appointment.
11. Turn D1310 code to "C" complete if counseling was completed.
12. Continue to counsel the patient at subsequent appointments as needed, making appropriate notes in DHOTEN.

PATIENT EDUCATION

The student is expected to:

1. Include patient education in every clinical appointment.
2. Perform a salivary flow rate assessment on each new patient appointment (prior to the pre-procedural rinse). Check the salivary pH. Record the results in the EHR
3. Complete the Limited Exam and analyze the findings relative to disease risk.
4. Disclose and record the O'Leary Plaque Index and discuss results with the patient showing them areas of biofilm.
5. Teach the appropriate toothbrushing technique and interdental plaque control and allow the patient to remove the existing biofilm (repeat this at each appointment). Record information in the EHR.
6. Complete the Oral Risk Assessment form and identify disease risks specific to the patient.
7. Recommend other preventive therapies relative to the risks identified and discuss these with the patient. Record in the EHR.
8. Explain disease processes to the patient in lay terms they can understand. Use illustrations, teaching aids and/or the microscope to enhance learning.
9. Demonstrate in the patient's mouth any OPT aids recommended and allow them to practice with feedback. Use communication and teaching principles and techniques such as modeling, prompting, cueing and feedback.
10. Select an appropriate treatment plan and explain it to the patient.

PERIODONTAL CHARTING

The student is expected to follow aseptic technique to:

1. Assemble mirror, Hu-Friedy PCP-UNC probe, and Nabers probe
2. Place the patient in the correct position.
3. Use light and mirror to aid in examination
4. Chart periodontal conditions of patients 18 years or older unless indicated by patient's condition.
5. Document in EHR the following:
 - a. Probe depths (PD measuring from base of the sulcus to the gingival margin)
 - b. Record six number readings per tooth. The measurement must be within 1mm of instructor's measurement
 - c. Bleeding on probing (BOP)
 - d. Free Gingival Margin (FGM measuring from gingival margin to CEJ). Areas of recessions are marked with negative numbers
 - e. Tooth mobility
 - f. Furcation involvement (using the Nabers probe)

PHASE-CONTRAST MICROSCOPE

The student is expected to:

Prepare the Equipment (if not already prepared)

1. Remove the dust cover. Do not remove the eyepiece lens covers. You will use the computer monitor to view the specimen.
2. Turn on the microscope light located at the back left side of the microscope base computer.
3. Turn on the microscope camera with the red button at the top of the microscope. When the microscope camera is on there will be illuminated red and blue lights.
4. Keep the light intensity control set to the lowest setting until needed.
5. Open the PixelPro program on the computer's Desktop.

End of Clinic

1. Turn off the microscope camera and light at the end of every clinic session.
2. Replace dust cover.
3. Place the slide in the biohazard sharp container.
4. Make the notation in EHR regarding the findings and information given the patient.
5. If you took photos, have faculty transfer photos to the patient's EHR before the end of the session. Photos left on the computer may be removed when IT runs computer updates.

Prepare the slide (Chair-side)

1. Explain the procedure to the patient and obtain consent.
2. Assemble the armamentarium:
 - a. glass slide
 - b. cover slip
 - c. sterile water in dropper dispensing bottle
 - d. scalers/curettes
3. Place one drop of water on the slide for each site to be examined.
4. Obtain the sample:
 - a. Make a clinical assessment of the mouth to determine the most advanced and severely diseased sites: deep crevices, adjacent to crowns, areas of malalignment.
 - b. Use a sterile instrument to obtain a sample from the most apical portion of the gingival crevice selected. A thin sample free of calculus is essential.
 - c. Use a second instrument to dislodge the sample into the drop of water. Do not break up the sample.
5. Place a single cover slip over the sample. Use an instrument to gently compress the cover slip.
6. Evaluate the sample. A thin sample free of calculus is essential.
7. Blot excess water with a tissue or paper towel.

View the Slide

6. Place the prepared slide with coverslip up onto the microscope stage.
7. Set the light intensity control knob to the highest setting.
8. Center the specimen over the light.
- 9. Use the computer monitor to view the specimen. Not the ocular lenses.**
10. Use the knobs under the stage to move the field into view.
 - a. The larger upper knob moves the slide front-to-back.
 - b. The smaller lower knob moves the slide left-to-right.
11. Rotate the objective turret until the 40X objective clicks into place above the specimen. The 40x objective is the best objective for observing the specimen and should not be changed.
12. Raise the stage: observe the distance between the objective and the cover slip of the slide.
13. Use the large outer knobs on either side of the microscope stand to raise the stage until the lens of the objective appears about to touch the cover slip.
14. Use the small fine focus knob (located outside of the knob to raise the stage) to bring the specimen into sharp focus.

15. During the clinic session, leave the camera on and return the light intensity control set to the lowest setting until needed.

Patient Education

1. Bring the patient to the microscope at this point.
2. Communicate information to the patient regarding the microorganisms present relative to their oral health.

Identify the Organisms Visible

State of Health

Some cocci
WBC's, 6/field or fewer

Some filamentous organisms
Low count vibrios

Microorganisms of marginal gingivitis

Cocci (TMC)
WBC's 0 – 12/field
Spinning and/or gliding rods
Trichomonads

Filamentous organisms
Spirochetes
Amoeba

Microorganisms of destructive periodontitis

Cocci (TNC)
WBC's
Spirochetal pumps
Gliding, palisading rods
Trichomonads

Filamentous organisms
Vibrios
Spirochetal brush forms
Amoeba

PIT AND FISSURE SEALANT APPLICATION

The student is expected to:

1. Review medical/dental history, general assessment, and oral inspection prior to treatment for information contraindicating treatment. Confirm treatment is planned by the student and approved by a dental faculty.
2. Explain procedure to patient and/or parent and get consent.
3. Assemble armamentarium:
 - mouth mirror
 - explorer
 - air/water syringe
 - articulating paper
 - articulating paper holder
 - cotton pellets or small sponges
 - etching material with tip
 - prime material (if applicable)
 - sealant material with application tip

- cotton rolls and Dri-angles and buccal shields
 - cotton tip applicator
 - pumice (moist)
 - microbrush
 - toothbrush or prophyl angle
 - slow handpiece
 - UV curing light with shield (from 3rd floor dispensary)
 - protective eyeglasses for operator and patient
 - floss
 - finishing burs or stones and high-speed handpiece
 - fluoride varnish
 - high speed suction
 - gauze
 - mouth prop (optional)
4. Position the patient comfortably. Utilize proper operator positioning.
 5. Place protective eyeglasses on the patient.
 6. Evaluate teeth scheduled for sealants with mirror and explorer.
 7. Mechanically cleanse the enamel with toothbrush, or prophyl angle brush and pumice.
 8. Thoroughly rinse the tooth surface with water.
 9. Isolate the tooth surface using cotton rolls and cotton roll holder and protect from any contamination.
 10. Dry the tooth surface with compressed air from 20-30 seconds.
 11. Follow the instructions of the desired sealant material.
 - a. Apply sealant to the etched surface. Be sure to use a yellow/orange light filter to prevent sealant material from curing too quickly.
 - b. Confine the sealant to the grooves and pits of applicable teeth by removing excess sealant material with a microbrush or placing the cotton tipped applicator distal to the tooth.
 - c. Have the patient close their eyes while visible light is on.
 - d. Place the end of the curing light tip (wand) 1-2 mm above the tooth surface.
 - e. Cure for the desired amount of time for sealant material 20-30 sec (according to manufacturers' instructions).
 - f. Gently Check for voids and retention of sealant with an explorer. Additional material can be added if the surface is not contaminated or wet.
 - g. Rinse with water or rub with a wet cotton roll to remove unpolymerized resin.
 - h. Check occlusion with articulating paper. If sealant material is too high, it can be adjusted using DH instruments, or a dentist can be notified to adjust the occlusion with a handpiece and bur. Occlusion must be rechecked with articulating paper. It is recommended that occlusion is checked when the patient is in supine and also sitting up to ensure an optimal and comfortable bite.
 - i. Floss between teeth after sealant placement.
 12. Have a supervising instructor check placement of sealant. If there is adjustment needed for the sealant, a dental faculty needs to be informed to correct sealant and the occlusion.
 13. Give fluoride treatment.
 14. Make complete, accurate, dated chart entries in EHR.
 15. Provide patient education material for the patient to take home.
 16. Maintain field:
 - a. Position light for maximum illumination.
 - b. Remove saliva and debris routinely to provide adequate vision and patient comfort.
 - c. Replace "wet cotton rolls".

Etching procedure:

1. Apply etchant for 15 – 60 seconds for permanent or deciduous teeth (see manufacturers' instructions for recommended etching time as it could differ depending on brand).

2. Apply etchant with a continuous, gentle dabbing motion.
3. Cover the grooves and pits with the etchant.
4. At least 2 – 3 mm of surrounding enamel around grooves and pits should be etched
5. Place the high-volume suction over the tooth
6. Rinse the tooth surface thoroughly with water for 60 seconds.
7. Dry the tooth surface with compressed air for 20 – 30 seconds.
8. The surface of the tooth should have a chalky white appearance, indicating complete etching.
9. The tooth should be isolated from the tongue, saliva, and tissue fluids using cotton rolls and/or dri-angles. Re-isolate with new cotton rolls or dri-angles.

Priming procedure (if applicable):

1. Apply PrimaDry and leave 5 seconds.
2. Dry by gently blowing the area with moisture-free and oil-free air.
3. Do not rinse.

POLISH/DEPLAQUE AND FLOSS

The student is expected to:

1. Assemble the armamentarium.
 - a. Check odontogram for types of restorations
 - b. Select correct abrasive agents
 - c. Use aseptic technique
2. Check the medical/dental history for information contraindicating the procedure.
3. Seat and position the patient comfortably. Utilize correct operator positioning.
4. Explain the procedure to the patient.
5. Discuss technique to be used with the instructor.
6. Place protective eyewear on the patient.
7. Inspect teeth for contraindications to polishing and select teeth to be polished.
8. Disclose the patient's mouth and discuss areas of plaque with the patient.
9. Use abrasive agents in order of most abrasive to least abrasive (if indicated) changing cups between abrasives.
10. Establish a fulcrum.
11. Use intermediate pressure & maintain slow, constant speed with the prophyl angle.
12. Flare the cup into the crevicular and proximal areas.
13. Adapt edge of cup to tooth contour.
14. Adapt occlusal brushes to pits and fissures.
15. Use auxiliary polishing aids as needed.
16. Use caution in retracting corners of the mouth or other soft tissue areas.
17. Wipe the cup clear of saliva and debris as needed to avoid splatter.
18. Floss all interproximal areas. (Refer to flossing procedure checklist).
19. Re-disclose and check with a mirror and air. Remove plaque as necessary.
20. Clean removable dentures or other removable appliances and return to the patient.
21. Evaluate the procedure and final product to determine ways to improve performance.
22. Clean up the treatment area and armamentarium.

POLISHING – SEATING POSITIONS

Operator Position	Area
FRONT	Max. Right Facial to Midline Max. Left Lingual to Canine
BEHIND	Max. Left Facial to Midline Max. Anterior Lingual Max. Right Lingual-Posterior
BEHIND	Mand. Left Facial to Midline Mand. Anterior Lingual- (Behind and Front)
FRONT	Mand. Right Facial to Midline Mand. Left Lingual-Posterior Mand. Right Lingual-Posterior

SCALING AND ROOT PLANING

The student is expected to:

1. Review patient assessment data to determine contraindications to treatment or other factors that will influence the procedure.
2. Assemble the appropriate armamentarium.
3. Use an aseptic technique.
4. Use correct patient and operator positioning.
5. Explain the procedure to the patient and present appropriate patient education and psychological support (i.e., homecare instructions, why the procedure is being done, postoperative instructions, etc.).
6. Evaluate the patient's pain and anxiety levels.
 - a. You may use Corah's Dental Anxiety scale to aid in this evaluation.
 - b. Use appropriate pain management techniques (topical Oraqix, and/or local anesthesia).
7. Review the patient assessment data and treatment plan assessment procedures to decide which area to scale, and consider the following variables:
 - a. Ability to complete the area within time available or required: Since partial calculus removal on a tooth is undesirable, select an area of the mouth that can be scaled and root planed to completion in the time available at this appointment.
 - b. Patient comfort and acceptance: To make the first scaling appointment less complicated and to help orient the patient to treatment, you may want to select either the sextant with the fewest teeth or the sextant with the least severe periodontal disease.
 - c. Need for tissue conditioning: Tissue conditioning is accomplished by initiating a daily program of plaque removal and warm salt water rinsing. The goals of such a program are:
 1. Gingival healing
 2. Lowered bacterial accumulation

3. Establishing plaque control behaviors by the patient.
 - d. Patient needs: When the patient indicates an area of discomfort, that area may be completed first.
 - e. Location: When two quadrants are to be treated at the same appointment, select a maxillary and mandibular quadrant on the same side of the mouth.
8. Formulate a plan as to the sequence of instruments to be used.
9. Use appropriate type, sharp, correctly contoured instruments according to the nature and location of the deposits.
10. Correctly grasp the instrument, establish fulcrum, position cutting edge and activate instrument according to the basic principles of instrumentation.
11. Completely scale each tooth so that:
 - a. all surfaces are calculus free
 - b. there is no undue tissue trauma.
12. Allow the patient to rinse thoroughly with water.
13. Evaluate the procedure and final product to determine that criteria were met and identify ways in which performance could be improved.
14. Make complete, accurate, dated chart entry into the EHR record.
15. Clean up the treatment area and armamentarium.

TOBACCO CESSATION COUNSELING & REFERRAL FOR ADDICTIONS

The student is expected to:

1. **Ask** every patient about tobacco or nicotine product use to identify all users. Determine if they have contemplated quitting.
2. **Advise** the tobacco/nicotine product user to quit by personalizing the message while utilizing active listening, sensitivity and empathy. Give information in a clear unambiguous way advising the patient to quit using tobacco. Make the message relevant to their periodontal disease, any identified pathology (tobacco melanosis, keratinization, nicotinic stomatitis, etc.), disease or medical history information (HBP, cardiovascular disease, diabetes, etc.).
3. **Assess** the patient's willingness to quit and level of addiction by administering the Tobacco History Form in the EHR. Assess the type, amount and frequency of use.

0-2 Very low dependence	6-7 High dependence
3-4 Low dependence	8-10 Very high dependence
5 Medium dependence	

4. Identify the stage of change and note this in the DHOTEN. Use Motivational Interviewing skills while going through the 5Rs by asking open ended questions, active listening, reflecting and summarizing (OARS).
5. **Assist** the patient with a quit plan, set a quit date (at least two weeks out) and make a note in the EHR. Discuss nicotine replacement and/or pharmacological therapies as appropriate for the patient's medical history and level of dependence. Refer to a Tobacco Dependence counselor or MD if more assistance or a prescription is needed.
6. **Follow-up** with a phone call in one week and document in the EHR. Follow-up at the next appointment with either encouragement or praise for quitting and follow the 5As and 5Rs.
7. **Add** D1320-Tobacco Counseling to the treatment plan selecting appropriate diagnostic code from the Harmful Oral Habits category. **Document** all activities performed in the treatment (T) and next appointment (N) sections of the DHOTEN note(s).
8. Demonstrate Professionalism

Examples of treatment notes from tobacco cessation counseling (should reflect activities performed and/or planned):

For Tobacco Cessation completed in one appointment with a pre-contemplative patient:

T: Assessed tobacco use and readiness: Moderately addicted and in the Precontemplation Stage.
Advised regarding personal risks and recommended patient quit.
N: 6-month recall. Revisit tobacco cessation readiness.

For tobacco cessation done in multiple appointments:Appt #1:

T: Assessed tobacco use and readiness: Severely addicted and in the planning stage. Advised regarding personal risks and recommended patient quit.
N: Discuss NRT and pharmacotherapy and set quit date

Appt #2:

T: Discussed Chantix and high dose nicotine patch. Pt. referred to MD. Quit date set for August 29th will begin Chantix 2 weeks prior.
N: Perio Re-eval. Follow-up on tobacco cessation.

Referral to Quitlines (a free service provided by the Department of State Health Services):

1. Patients can receive up to 5 counseling sessions and a 2-week supply of over-the-counter NRT. Following the referral, the Quitline proactively calls the patient within approximately 48 hours to schedule the counseling sessions; once the patient enrolls in the service, the NRT is mailed to their home (TX, USA address)
 - a. Patient who are 18 or older receive the full range of services
 - b. Patients who are 13-17 can receive counseling only
2. Quitline services are available 24/7 and counseling is provided in the patient's language of choice
3. Referral tools:
 - EMR referral: with the eTobacco Protocol integrated in the EMR, there can be an added button located near the question regarding the "tobacco use" status that creates an electronic referral to the Quitline with the patient's approval.
 - Referral by apps: Help to Quit and Texas Quitline apps are available for Apple and Android devices (community health worker outreach, health fairs for schools, school-based clinics).
 - Fax: referral by fax with a printable form available here: www.yesquit.org

For suspected alcohol and drug abusers refer to:

Alcoholics Anonymous (AA) <https://www.aa.org/> multiple locations in Houston area
Narcotics Anonymous (NA) <https://www.na.org/meetingsearch/> multiple locations in Houston area

For suspected marijuana users:

Resources for providers <https://www.cdc.gov/marijuana/index.htm>
National Institute on Drug Abuse <http://www.drugabuse.gov/drugs-abuse/marijuana>
National Council on Alcoholism and Drug Dependence <https://ncadd.org/>
American Academy of Family Physicians <http://www.aafp.org/about/policies/all/marijuana.html>
MedlinePlus <https://www.nlm.nih.gov/medlineplus/marijuana.html>
National Cancer Institute <http://www.cancer.gov/about-cancer/treatment/cam/hp/cannabis-pdq>

ORAL HYGIENE INSTRUCTIONS: TOOTHBRUSHING

The student is expected to:

1. Assess patient needs: medical history, oral exam, gingival description, case classification.
2. Determine oral hygiene regimen and nutritional habits.
3. Question as to when dental home care procedures were last performed.
4. Disclose the patient.
5. Complete and record the plaque index.
6. Provide brushing instructions:
 - a. Allow the patient to brush in the usual way.
 - b. Rediscover. Request patient to identify remaining plaque deposits. The patient should be seated upright in the dental chair with the light and hand mirror angled to aid their visualization.
 - c. Suggest alteration in brushing technique, if appropriate.
 - d. Demonstrate brushing in the patient's mouth. Request the patient to perform in their own mouth. Make modifications, as necessary, and incorporate aspects of other techniques, as appropriate.
 - e. Give constructive feedback and reinforcement.
7. Demonstrate tongue brushing.
8. Have patient:
 - a. Extend the tongue
 - b. Place the brush as far posteriorly as possible
 - c. Sweep the brush anteriorly, displacing the tongue as little as possible.

Bass Technique of Brushing

Facial and Lingual Surfaces**

1. Grasp the toothbrush in order to maintain control during all movements.
2. Point the bristles apically at a 45° angle to the long axis of the tooth.
3. Place the bristles at the gingival margin. The first row of bristles will be close to the crevice. The adjacent row will touch the gingival margin.
4. Press lightly. The bristles will contour themselves into the crevice and interproximal area.
5. Apply 10 short back-and-forth vibratory or circular strokes. Do not lift the brush or use a scrubbing motion.
6. Relax the bristle pressure and move the brush to the next segment, overlapping at least one tooth.

Anterior Lingual Surfaces**

1. Insert the brush vertically.
2. Place the bristles of the toe of the brush at the crevicular area and vibrate.
3. Pull the bristles over the tooth surface toward the incisal edge.

Occlusal Surfaces**

1. Scrub by moving the bristles back and forth

***Follow a definite sequence

TREATMENT PLAN/REFUSAL OF TREATMENT

The student is expected to:

1. Complete all patient records and diagnostic aids, assessing all information which will influence dental hygiene treatment. Forms and procedures include medical and dental history, extra/intraoral exam, gingival description, dental charting, periodontal charting, and oral risk assessment.
2. Using all patient assessment data, determine a dental hygiene diagnosis. In periodontally involved cases, the dental hygiene diagnosis should include type, extent, and severity of periodontal disease.

3. Based on dental hygiene diagnosis and other assessment data, prepare a comprehensive treatment plan for patient care, including dental hygiene therapeutic services and other preventive services as determined from assessment data.
4. Enter a sequential treatment plan into the EHR.
5. Discuss the comprehensive treatment plan with the instructor and patient BEFORE having the instructor approve. Informed consent includes presenting the treatment options, discussing alternatives and risks, and allowing the patient to ask questions before signing the treatment plan.
6. If a patient refuses any part of the treatment plan (including fluoride or radiographs), a Refusal of Treatment Form, must be completed and signed by the patient, student, DH faculty and DDS faculty. This form should be approved by the DDS faculty, printed and returned to the patient. The student must document this in the DHOTEN. Students can create a "copy" of the treatment under the Treatment Planning module and have the DH faculty approve the second option without the treatment that the patient refused. This way, it documents that the procedure was recommended and presented, but declined by the patient.
7. Once the comprehensive treatment plan is signed by the patient, make an entry in DHOTEN that the treatment plan has been discussed with the patient, and informed consent was signed on _____ (date).
8. Make complete, accurate, chart entry in the EHR.
9. Demonstrate Professionalism.

ULTRASONIC SCALERS

The student is expected to:

1. Review medical/dental history, vital signs, chart, and patient assessment form for data that contraindicates proceeding with treatment or will otherwise influence the procedure. Patients with a pacemaker should have the Piezo used, not the magnetostrictive Cavitron.
2. Explain the procedure and rationale to the patient, providing individualized patient education.
3. Have the patient use a pre-procedural rinse before using the ultrasonic scaler.
4. Assemble armamentarium:
 - a. ultrasonic scaling unit
 - b. plastic ultrasonic drape for patient
 - c. pre-procedural rinse
 - d. glasses for patient
 - e. paper towels
 - f. face mask & shield
 - g. surgical cap/bonnet
 - h. earplugs
 - i. mouth mirror
 - j. saliva ejector and HVE
 - k. ultrasonic scaler inserts
5. Connect the BLUE water line to the water outlet. Make sure to turn on the water connector knob at the attachment port.
6. Make sure that the ultrasonic unit is securely plugged in and turn on the unit (Cavitron only).
7. Make sure HVE suction is ready to use.
8. "Bleed" ultrasonic water line for two minutes.
9. Check ultrasonic insert for damage.
10. Fill the ultrasonic handpiece with water and insert appropriate insert. If using the Piezo, attach the appropriate Piezo tip to the unit handpiece.
11. Drape the patient with a plastic apron and provide tissues and safety glasses.
12. Make sure to have a mask, protective eyewear, face shield, earplugs, surgical cap/bonnet, and gloves on. Follow aseptic technique and infection control protocol.
13. Use the appropriate insert for the calculus and/or plaque that is present in the patient's mouth.

14. Adjust the ultrasonic unit to the correct power setting and water flow for the tip that you are using.
15. Use correct handpiece cord management.
16. Adjust patient position to the proper angle.
17. Using a modified grasp, with light touch and appropriate fulcrum
18. Apply the instrument to the teeth using correct angulations.
 - a. Piezo tip- utilizes lateral surfaces and lower 2 mm of tip
 - b. Ultrasonic- utilizes lower 2 mm of tip
19. Keep the working end in constant motion, controlled, and short overlapping strokes.
20. Maintain visibility by keeping the correct patient/operator position.
21. Scale the teeth utilizing a system that minimizes trauma to both teeth and gingival tissue, changing to appropriate ultrasonic inserts as necessary in order to adapt to changing tooth topology, and filling the handpiece with water before inserting another ultrasonic insert.
22. Use HVE suction continuously.
23. Check for patient comfort both verbally and visually.
24. Give pre-and post-operative instructions.
25. Assess procedures and outcomes and determine ways to improve performance.
26. Clean and disinfect cubicle and ultrasonic unit.
27. Rinse and dry ultrasonic insert/s, place in sterilization cassette and place on cart for sterilization.
28. Make complete, accurate, dated chart entries in EHR.

VITAL SIGNS

The student is expected to:

1. Explain the procedure and rationale to the patient.

Take Blood Pressure

2. Have the patient sit upright, roll up sleeve, flex arm slightly and rest on the arm of the chair at heart level and legs uncrossed.
3. Disinfect ear pieces to the stethoscope.
4. Put a sphygmomanometer cuff one inch above the elbow with a compression bag over the brachial artery.
5. Place a manometer where it can be easily read.
6. Palpate the radial artery and inflate the cuff until the pulse disappears, continue to inflate another 20-30 mmHg. The cuff should be deflated slowly at 2-3 mm HG per second until the radial pulse reappears. The point at which the pulse disappears and then reappears on deflation is called palpatory systolic pressure. Release all pressure in the cuff.
7. Place the stethoscope over the brachial artery slightly below the cuff and inflate the cuff to 20-30 mmHg. above the previously determined palpatory systolic pressure.
8. Release the pressure at a rate of 2-3 mmHg/second.
9. Listen for systolic and note the number. Listen for diastolic and note the number.
10. Record the systolic over diastolic pressure, e.g. 120/80 in progress notes.
11. Recognize abnormally high readings.
12. Repeat procedure if reading is not within the normal range. (**Located on the UTSD intranet <https://inside.uth.edu/dentistry/docs/mgmtofthehypertensivepatientpolicy.pdf>).**
13. Tactfully advise the patient about the reading.
14. Note: For a patient with blood pressure significantly above normal ($\geq 180^*$ or $\geq 120^*$) advise the faculty and have the Dispensary RN recheck the readings.

Take the Pulse

15. Palpate for 60 seconds a readily available artery –usually either the radial or brachial artery. Press the fleshy portion of the index and middle fingers onto the patient's skin gently enough to feel the pulsation but not so firmly that the pressure occludes the artery. Use the carotid artery, if the radial pulse is not able to be detected.-
16. Evaluate and record the rate, e.g. 60/min. in progress notes of the patient's chart.
17. Advise the faculty if reading exceeds 80 beats per minute.

Measure Respiration

18. Observe the patient's chest unobtrusively by keeping the fingers on the patient's pulse as if continuing to take the pulse.
19. Observe the rise and fall of the chest for a minimum of 30 seconds (ideally for one minute).
20. Measure the rate of respiration and record it accurately, e.g. 14/min. in progress notes.
21. Advise the faculty if readings diminish to 12 per minute or exceed 28 per minute.

Take Temperature

1. Using a Temp-a-Dot found at the dispensary, peel back the wrapper to expose the handle end of the thermometer.
2. Remove the thermometer taking care not to touch that part which is placed in the patient's mouth. Place the thermometer under the tongue as far back as possible into either heat pocket.
3. Have the patient press tongue down on the thermometer, keeping mouth closed.
4. Keep the thermometer in the patient's mouth for one minute.
5. After removal, allow ten seconds before reading the temperature. The last blue dot indicates temperature.
6. Record temperature in progress notes of patient chart under H in the DHOTEN.
7. Evaluate the patient's vital signs and correlate them with other physical and medical history findings to determine the treatment plan.